



Bookmarks

- ▶ [Module 1: The Basics of R and Introduction to the Course](#)
- ▶ [Entrance Survey](#)
- ▶ [Module 2: Fundamentals of Probability, Random Variables, Distributions, and Joint Distributions](#)
- ▶ [Module 3: Gathering and Collecting Data, Ethics, and Kernel Density Estimates](#)
- ▶ [Module 4: Joint, Marginal, and Conditional Distributions & Functions of Random Variable](#)

Module 9: Single and Multivariate Linear Models > The Linear Model > Goodness of Fit - Quiz

Goodness of Fit - Quiz

Bookmark this page

Question 1

1/1 point (graded)

True or False: The formula for $R^2 = SSR/SST$.☐ a. True☒ b. False ✓

Explanation

The formula for R^2 is $1 - SSR/SST$. We define R^2 with the "1 minus" out in front so that a larger R^2 means that the fit is better (that more of the variation in Y is explained by variation in X).

You have used 1 of 1 attempt

✓ Correct (1/1 point)

- ▶ [Module 5: Moments of a Random Variable, Applications to Auctions, & Intro to Regression](#)
- ▶ [Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation](#)
- ▶ [Module 7: Assessing and Deriving Estimators - Confidence Intervals, and Hypothesis Testing](#)
- ▶ [Module 8: Causality, Analyzing Randomized Experiments, & Nonparametric Regression](#)
- ▼ [Module 9: Single and Multivariate Linear Models](#)

The Linear Model

due Nov 28, 2016 05:00 IST



Question 2

1/1 point (graded)

$$(n - 2) \frac{R^2}{1 - R^2}$$

The expression above has an F-distribution under the null hypothesis. We use the expression above to test that hypothesis that:

☐ a. $\beta_1 \neq 0, \dots, \beta_k \neq 0$

☐ b. $\beta_1 \neq \dots \neq \beta_k$

☐ c. $\beta_1 = \dots = \beta_k$

☒ d. $\beta_1 = \dots = \beta_k = 0$ ✓

Explanation

In addition to using R^2 as a basic measure of goodness-of-fit, we can also use R^2 as the basis of a test of the hypothesis that our coefficients are all zero. (This would mean that our regressors do not explain our dependent variable.)

The Multivariate Linear Model

due Nov 28, 2016 05:00 IST



Module 9: Homework

due Nov 21, 2016 05:00 IST



- ▶ Module 10: Practical Issues in Running Regressions, and Omitted Variable Bias

- ▶ Exit Survey

Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)

Question 3

1/1 point (graded)

We should reject the hypothesis from question (2) when the expression above is:

☐ a. Small

☒ b. Large ✓

Explanation

The expression above will be large when R^2 is large. R^2 is large when our SSR is much smaller than our SST , meaning that the variation in \mathbf{X} explains a lot of the variation in \mathbf{Y} . This would suggest that the coefficients β on our \mathbf{X} 's are non-zero with high probability.

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Discussion

Topic: Module 9 / Goodness of Fit - Quiz

[Show Discussion](#)

© All Rights Reserved



© 2016 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

