



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

44:55:04



< Previous



## 11. Computing partial derivatives

Bookmark this page



Calculator




Hide Notes



Explore

Examples

[Start of transcript. Skip to the end.](#)



PROFESSOR: So how to compute these things?

Oh, there's a piece of notation I haven't told you yet.

So another notation you will see,

I think this is what one uses a lot in physics

and this is what one uses a lot in applied math, which

is the same thing as physics but with different notations.



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

To compute the partial derivative of  $f(x, y)$  with respect to  $x$ , we treat  $y$  as a constant and differentiate each term with respect to  $x$  only. To illustrate this, let's first think about derivatives in the single variable case.

In order to compute

$$\frac{d}{dx} \sin(7x)$$

we use the chain rule to obtain

$$\frac{d}{dx} \sin(7x) = \cos(7x) \cdot \frac{d}{dx}(7x) = 7 \cos(7x).$$

Now, let's say we want to compute

$$\frac{\partial}{\partial x} \sin(yx).$$

To do this, we use the chain rule as we did above, except now  $y$  is playing the role of  $7$ . So we compute

$$\frac{\partial}{\partial x} \sin(yx) = \cos(yx) \cdot \frac{\partial}{\partial x}(yx) = y \cos(yx).$$



Calculator



Hide Notes

This is what we mean when we say to treat  $y$  as a constant. Similarly, to compute the partial derivative of a function  $f(x, y)$  with respect to  $y$ , we treat  $x$  as a constant and differentiate each term with respect to  $y$  only.

Let's do some worked examples before you get the chance to practice on your own.

**Example 11.1** Consider  $f(x, y) = x^2 + 3xy$ . Then

$$f_x(x, y) = \frac{\partial}{\partial x}(x^2 + 3xy)$$

(2.12)

$$= \frac{\partial}{\partial x}x^2 + 3y\frac{\partial}{\partial x}x$$

(2.13)

$$= 2x + 3y$$

(2.14)

and

$$f_y(x, y) = \frac{\partial}{\partial y}(x^2 + 3xy)$$

(2.15)

$$= \frac{\partial}{\partial y}x^2 + 3x\frac{\partial}{\partial y}y$$

(2.16)

$$= 0 + 3x$$

(2.17)

$$= 3x.$$

(2.18)

**Example 11.2** Let  $g(x, t) = \sin(x - 10t)$ . Then

$$g_t(x, t) = \frac{\partial}{\partial t}\sin(x - 10t)$$

(2.19)

$$= \cos(x - 10t) \cdot \frac{\partial}{\partial t}(x - 10t)$$

(2.20)

$$= -10 \cos(x - 10t)$$

(2.21)

and

$$g_x(x, t) = \frac{\partial}{\partial x}\sin(x - 10t)$$

(2.22)

$$= \cos(x - 10t) \cdot \frac{\partial}{\partial x}(x - 10t)$$

(2.23)

$$= \cos(x - 10t).$$

(2.24)

11. Computing partial derivatives

Hide Discussion

**Topic:** Unit 1: Functions of two variables / 11. Computing partial derivatives

Add a Post

Show all posts

by recent activity

There are no posts in this topic yet.

◀ Previous

Next ▶

© All Rights Reserved



## edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

## Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

## Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)



© 2021 edX Inc. All rights reserved.  
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)

Calculator

Hide Notes