



### 9.3.4 4th order Runge-Kutta Method (RK4)

Bookmark this page

MO2.7

The most popular form of a fourth-order ( $p = 4$ ) accurate Runge-Kutta method is:

$$\underline{a} = \Delta t \underline{f}(\underline{v}^n, t^n) \tag{9.10}$$

$$\underline{b} = \Delta t \underline{f}(\underline{v}^n + \underline{a}/2, t^n + \Delta t/2) \tag{9.11}$$

$$\underline{c} = \Delta t \underline{f}(\underline{v}^n + \underline{b}/2, t^n + \Delta t/2) \tag{9.12}$$

$$\underline{d} = \Delta t \underline{f}(\underline{v}^n + \underline{c}, t^n + \Delta t) \tag{9.13}$$

$$\underline{v}^{n+1} = \underline{v}^n + \frac{1}{6}(\underline{a} + 2\underline{b} + 2\underline{c} + \underline{d}) \tag{9.14}$$

Note that this method requires four evaluations of  $\underline{f}$

< Previous

Next >

### Discussions

All posts sorted by recent activity



# edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

## Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

## Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.  
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)