



[Course](#) > [Unit 1: Fourier Series](#) > [1. Introduction to Fourier Series](#) > 11. A worked example

**Audit Access Expires Jun 24, 2020**

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 11. A worked example


Find the Fourier series of the  $2\pi$ -periodic function

$$f(t) = \begin{cases} t & 0 < t < \pi \\ 0 & -\pi < t < 0 \end{cases}.$$

**Solution:** In this example, we will work out the terms  $a_n$  and leave it to you to find the coefficients  $b_n$ .

First, draw a picture.





Next, find the constant term:

$$\frac{a_0}{2} = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) dt = \frac{\text{area under one period of curve}}{\text{length of period}} = \frac{\pi^2/2}{2\pi} = \frac{\pi}{4}.$$

Now for the other terms  $a_n$ :

$$\begin{aligned} a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \cos(nt) dt \\ &= \frac{1}{\pi} \int_{-\pi}^0 0 \cos(nt) dt + \frac{1}{\pi} \int_0^{\pi} t \cos(nt) dt \\ &= \frac{1}{\pi} \int_0^{\pi} t \cos(nt) dt \end{aligned}$$

Evaluating this integral, we find

$$\frac{1}{\pi} \int_0^{\pi} t \cos(nt) dt = \frac{1}{\pi} \frac{\cos(n\pi) - 1}{n^2} = \frac{1}{\pi} \frac{(-1)^n - 1}{n^2} = \begin{cases} \frac{-2}{\pi n^2} & n \text{ odd} \\ 0 & n \text{ even} \end{cases}.$$

Therefore the Fourier series is given by



$$f(t) = \frac{\pi}{4} + \sum_{n \text{ odd}} \frac{-2}{\pi n^2} \cos(nt) + \sum_n b_n \sin(nt).$$

We leave the computation of the  $b_n$  as an exercise for you.

## Finish the example

1/1 point (graded)

Find the coefficients  $b_n$  of the Fourier series for the  $2\pi$ -periodic function

$$f(t) = \begin{cases} t & 0 < t < \pi \\ 0 & -\pi < t < 0 \end{cases}.$$

(Find a formula that holds for both  $n$  even and  $n$  odd.)

$b_n =$

✓ Answer:  $(-1)^{(n+1)}/n$

### Solution:

We can use computer assistance or integration by parts to see that

$$b_n = \frac{1}{\pi} \int_0^\pi t \sin(nt) dt = \frac{-\cos(n\pi)}{n} = \frac{(-1)^{n+1}}{n}.$$

Submit

You have used 2 of 5 attempts



## 11. A worked example

**Topic:** Unit 1: Fourier Series / 11. A worked example

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Learn About Verified Certificates

© All Rights Reserved

