<u>Help</u>

sandipan_dey >

<u>Calendar</u> **Discussion Notes** <u>Course</u> <u>Progress</u> <u>Dates</u>

☆ Course / Unit 2: Geometry of Derivat... / Lecture 4: Introduction to vectors and dot pro...



You are taking "Exam (Timed, No Correctness Feedback)" as a timed exam. Show more

End My Exam

Previous

44:22:20





☐ Bookmark this page



Explore

Vectors in 2D



PROFESSOR: So we draw a vector as

Start of transcript. Skip to the end.

We'll say this is the vector v. And we can describe v

with numbers.

So this here is called v1.

And this here is called v2.

And then the vector v is going to be written [v1, v2].

0:00 / 0:00

66 ▶ 2.0x X CC

Video

Download video file


Transcripts

Download SubRip (.srt) file Download Text (.txt) file

For now, we'll work with 2-dimensional vectors. Later in the course, we will discuss 3-dimensional and higher dimensional vectors. We will have some spoilers about these ideas in case you are interested now.

Example 3.1

The vector $ec{v} = \langle -1, 5
angle$ is represented by an arrow starting at the origin and extending to the point in the coordinate plane given by (x,y)=(-1,5). The image below shows the vector in blue. The lengths of its components are indicated by the dashed orange lines.




■ Calculator

Example 3.2

Sometimes the components of a vector are described using variables or parameters. For example, a vector

$$ec{v} = \langle v_1, v_2
angle$$

is shown below.



→ Spoiler: vectors in 3D

A vector in three dimensions has three components and can be written as

$$\vec{v} = \langle v_1, v_2, v_3 \rangle$$
 (3.1)

or

$$\vec{v} = v_1 \,\hat{i} + v_2 \,\hat{j} + v_3 \,\hat{k} \tag{3.2}$$

where $\hat{\pmb{i}}$, $\hat{\pmb{j}}$, and $\hat{\pmb{k}}$ are the unit vectors in the \pmb{x} , \pmb{y} , and \pmb{z} directions, respectively. In other words,

$$\hat{i} = \langle 1, 0, 0 \rangle$$

$$\hat{j} = \langle 0, 1, 0
angle$$


$$\hat{k} = \langle 0, 0, 1 \rangle.$$

Example 3.3

The vector $ec{v} = \langle 1, 2, 3 \rangle$ is represented by an arrow starting at the origin and extending to the point in the coordinate system given by (x,y,z)=(1,2,3) . The image below shows the vector in blue. The lengths of its components are indicated by the dashed orange lines.

► VECTOR IN 3 DIMENSIONS





A note about notation:

In this course, Professor Larry Guth uses the notation

$$ec{v}=(v_1,v_2)$$
 .

In this text and in videos with Professor Denis Auroux, we use the notation

$$ec{v} = \langle v_1, v_2
angle$$

to clarify any ambiguity between when we are talking about points or vectors.

Other common notation:

$$ec{v} = \langle v_1, v_2
angle$$

$$ec{v} = [v_1, v_2]$$

$$ec{v} = egin{pmatrix} v_1 \ v_2 \end{pmatrix}$$


$$ec{v} = v_1 \hat{i} + v_2 \hat{j},$$

where $\hat{\mathbf{i}}$ and $\hat{\mathbf{j}}$ are the unit vectors in the x and y directions, respectively. In other words,

$$\hat{i} = \langle 1, 0 \rangle$$

$$\hat{j} = \langle 0, 1
angle.$$

The vector \vec{v} in the above example was drawn starting at the origin. But vectors do not have to start at the origin.



In the above picture,

$$ec{v} = \langle -1, 1
angle$$

where the -1 represents the change in the $m{x}$ coordinate and the $m{1}$ represents the change in the $m{y}$ coordinate. We also have

$$ec{w} = \langle -1, 1
angle = ec{v}.$$

Two vectors are equal if they have the same magnitude (or length) and point in the same direction.

3. Vector examples in 2 dimensions

Hide Discussion

Topic: Unit 2: Geometry of Derivatives / 3. Vector examples in 2 dimensions

Add a Post

Show all posts \checkmark by recent activity \checkmark		
Q	Voice almost inaudible In my laptop, in professor Guth videos, his voice is almos inaudible at top sound settings. This is not the case with other videos of th	5
?	[Staff] Small typo In the text in "other common notation" section should be "in the x and y directions" instead of "in the xand y directions". Space need	2
Q	Subtitle Mistake The subtitles say > What does it mean for two factors to be equal to each other? When it should say >What does it mean for two *ve	2

Previous	Next >
Previous	Next >

© All Rights Reserved



edX

About

Affiliates

edX for Business

Open edX

Careers

<u>News</u>

Legai

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

Trademark Policy

<u>Sitemap</u>

Connect

<u>Blog</u>

Contact Us

Help Center

Media Kit

Donate















© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>