

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



MITx: 6.86x

Machine Learning with Python-From Linear Models to Deep Learning

[Help](#)[sandipan_dey](#)

[Unit 2 Nonlinear Classification,](#)
[Linear regression, Collaborative](#)
[Course](#) > [Filtering \(2 weeks\)](#)

> [Project 2: Digit recognition \(Part 1\)](#) > 5. Temperature

5. Temperature

We will now explore the effects of the temperature parameter in our algorithm.

You will be working in the files `part1/main.py` and `part1/softmax.py` in this problem

Effects of Adjusting Temperature

1/1 point (graded)

Explain how the temperature parameter affects the probability of a sample $x^{(i)}$ being assigned a label that has a large θ . What about a small θ ?

- ☐ Larger temperature leads to less variance
- ☒ Smaller temperature leads to less variance
- ☐ Smaller temperature makes the distribution more uniform



You have used 3 of 3 attempts

✓ Correct (1/1 point)

Reporting Error Rates

2.0/2.0 points (graded)
Set the temperature parameter to be 0.5, 1, and 2; re-run `run_softmax_on_MNIST` for each one of these (add your code to the specified part in `main.py`).

Error_{|T=0.5} = ✓

Error_{|T=1} = ✓

Error_{|T=2} = ✓

Submit

You have used 1 of 20 attempts

Return the temp_parameter to be 1 before moving on to the next section.

Discussion

Hide Discussion

Topic: Unit 2 Nonlinear Classification, Linear regression, Collaborative Filtering (2 weeks):Project 2: Digit recognition (Part 1) / 5. Temperature

Add a Post

◀ All Posts

What is "Effects of Adjusting Temperature" question asking

question posted 4 days ago by **CoolZ** (Community TA)

Anybody get what this question is asking?

Basically higher temperature will make all outcomes more equiprobable and lower temperature causes a greater difference. But which distribution is the question refer to?

This post is visible to everyone.

Mark B2 (Community TA)

4 days ago - marked as answer 4 days ago by **CoolZ** (Community TA)

$$p(y(i) = j|x(i), \theta)$$



...

Thanks, that means "variance" in the question is not refer to this distribution's variance. Then it must be the variance of probability of a sample $x^{(i)}$ being assigned a label that has a large θ .

This feels more like a word puzzle then a ML question. 😞

posted 4 days ago by [Cool7](#) (Community TA)

...

""variance" in the question is not refer to this distribution's variance."
It does. It returned by compute_probabilities function and temperature is one of parameters. Would temperature change the returned variance?

posted 4 days ago by [Mark B2](#) (Community TA)

...

Oh! I don't know what was I thinking, confused myself. I really should stop doing things at night,embarrassing. Thanks very much.

posted 4 days ago by [Cool7](#) (Community TA)

...

I'm definitely confused on this one. If a higher temperature makes all outcomes more equiprobable then why is a decrease in variance for larger temperature apparently not correct?

posted 4 days ago by [synnfusion](#)

...

The variance of uniform distribution on [0, 1] is 1/12, the variance of delta function is zero.

posted 3 days ago by [Mark B2](#) (Community TA)

...

Might be helpful if I remember that variance depends on distance from mean and not how much the pdf is going up and down.

posted 3 days ago by [synnfusion](#)

Add a comment

Add a Response


3 other responses

[sandipan dey.](#)
2 days ago

+

...

Already used 2 attempts and yet to get the correct answer. Used a random theta matrix and plotted the class-prob-distribution and temperature-vs-variance with the probabilities obtained using compute_probabilities() ,for a test datapoint. The plot supports the intuition, although the grader does not accept the answer and i am not sure what i am doing wrong. Any help? Thanks in advance.



I think from the earlier comments, this question is asking a different thing, to what I think it's asking!

posted 2 days ago by [robweatherston18](#)

...

Check your variance function, note that np.var is not appropriate to calculate variance in this case.

posted about 23 hours ago by [nontawat](#)

...

Why is np.var inappropriate @nontawat? I thought the question is asking the variance of the computed probabilities (one for each class). Even if we use ddof=1, the result will not change.

posted about 14 hours ago by [sandipan_dey](#).

...

Hint: look at your plot and figure out which kind of distribution it is.

posted about 6 hours ago by [nontawat](#)

Add a comment

[sakimarquis](#)
a day ago

+
...

same here, i use all 3 attempts just guessing, it's really confusing.

Add a comment

[dkhachatrian](#)
about 14 hours ago

+
...

What @Mark_B2 described is what the grader is asking for. Staff should probably make this more explicit. Especially because usually, looking at the distribution of the *ID number corresponding to a predicted label for a classification problem* doesn't have much meaning. For example, what if the label-encoding were {0: "cat", 1: "dog", 2:"other"} rather than our current example of {0:'0', 1:'1', ..., 9:'9'}? (I imagine a more "meaningful" distribution to look at could be the 1-D distribution of probabilities -- to see whether/how often the classifier is "confident" vs "ambivalent".)

Add a comment

Showing all responses

Add a response:

Preview

Submit

Learn About Verified Certificates