



< Previous



Next >

## 5. Justification

🔖 Bookmark this page



Calculator




Hide Notes



Explore

Justify the Chain Rule

[Start of transcript. Skip to the end.](#)



PROFESSOR: I mean, you probably have a right to expect some reason for why this works. Why is this valid? After all, I first told you we have this new mysterious object. And then I'm telling you, well, we can do that. But, you know, I kind of have pulled it

▶

0:00 / 0:00

▶ 2.0x

🔊

🔍

CC

“

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)  
[Download Text \(.txt\) file](#)

From now on, we will sometimes just use the phrase "chain rule" instead of "multivariable chain rule."

Why is the chain rule true?

On the previous page, we claimed that you could “divide everything by  $dt$ ” to obtain the chain rule. To justify this requires clearly interpreting differentials. However, the justifications given here are not entirely rigorous, and are just meant to give you an idea of what is going on when we write equations with differentials.

1st attempt

One justification is that, if  $x, y$ , and  $z$  are all functions of  $t$ , then we can compute  $dx, dy$ , and  $dz$  directly using single-variable calculus.

$$dx = x'(t) dt, \quad dy = y'(t) dt, \quad dz = z'(t) dt.$$

(6.128)

By substitution, we obtain the "chain rule" statement made on the previous page.

$$df = f_x x'(t) dt + f_y y'(t) dt + f_z z'(t) dt$$

(6.129)

$$= (f_x x'(t) + f_y y'(t) + f_z z'(t)) dt$$

(6.130)

Now we have an equation for  $df$  in terms of only  $t$  and  $dt$ . The coefficient on  $dt$  must be the derivative of  $f$  with respect to  $t$ .

2nd attempt

🧮

Calculator

🖋️

Hide Notes

Another way to convince ourselves of the chain rule is to replace the  $d$ 's by  $\Delta$ 's. This removes any ambiguity about the meaning.

We know from linear approximation that

$$\Delta f \approx f_x \Delta x + f_y \Delta y + f_z \Delta z \tag{6.131}$$

Dividing by  $\Delta t$  gives


$$\frac{\Delta f}{\Delta t} \approx \frac{f_x \Delta x + f_y \Delta y + f_z \Delta z}{\Delta t} \tag{6.132}$$

$$\frac{\Delta f}{\Delta t} \approx f_x \frac{\Delta x}{\Delta t} + f_y \frac{\Delta y}{\Delta t} + f_z \frac{\Delta z}{\Delta t} \tag{6.133}$$

Now if we imagine  $\Delta t$  moving towards zero, we can replace all  $\Delta$ 's with  $d$ 's to get a correct statement.

Example of the Chain Rule

Start of transcript. Skip to the end.



0:00 / 0:00

2.0x

🔊

🔍

📄

🗣️

PROFESSOR: OK, so let's check maybe things in an example.

Let's say that we really don't have any faith in these things.

And so let's try to do it.

So let's say I give you a function that's  $x$  squared  $y$  plus  $z$ .

And let's say that maybe  $x$  will be  $t$ ,  $y$  will be  $e$  to the  $t$ , and  $z$  will be  $\sin t$ .

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)  
[Download Text \(.txt\) file](#)

**Example 5.1** Let's look at an example of the chain rule in practice. Suppose we have a quantity  $w$  that depends on  $x$ ,  $y$ , and  $z$  as

$$w = x^2 y + z \tag{6.134}$$

Now suppose we can't control  $x$ ,  $y$ , and  $z$  directly, but they each depend on the parameter  $t$ :

$$x(t) = t \tag{6.135}$$

$$y(t) = e^t \tag{6.136}$$

$$z(t) = \sin t \tag{6.137}$$

Changing the variable  $t$  will cause  $w$  to change, and we would like to know the corresponding rate of change  $\frac{dw}{dt}$ .

The chain rule tells us that

$$\frac{dw}{dt} = \underbrace{(2xy)}_{w_x} \frac{dx}{dt} + \underbrace{x^2}_{w_y} \frac{dy}{dt} + \underbrace{1}_{w_z} \frac{dz}{dt}$$

(6.138)

Now substituting in the formulas for  $x$ ,  $y$ , and  $z$  gives a final answer:

$$\frac{dw}{dt} = 2te^t + t^2e^t + \cos t$$

(6.139)

Checking the answer with the old way

Since the above method used a new technique (the chain rule) let's make sure we get the same answer using our old methods. Namely, in this case, we can write down the full formula for  $w$  as a function of  $t$  and take a single-variable derivative.

$$w = x^2y + z$$

(6.140)

$$w(t) = t^2e^t + \sin t$$

(6.141)

$$w'(t) = 2te^t + t^2e^t + \cos t$$

(6.142)

Indeed, the answers match.

5. Justification

Hide Discussion

Topic: Unit 5: Curves and Surfaces / 5. Justification

Add a Post

Show all posts ▾by recent activity ▾

There are no posts in this topic yet.

✕

< Previous

Next >



Calculator

Hide Notes

# edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

## Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

## Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)



© 2021 edX Inc. All rights reserved.  
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)