



[Course](#) > [Unit 2:...](#) > [4 Eigen...](#) > 16. Ap...

16. Applications of diagonalization

Powers of a matrix



7:37 / 7:37



2.0x



HD



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Suppose that $\mathbf{A} = \mathbf{S}\mathbf{D}\mathbf{S}^{-1}$. Then

$$\mathbf{A}^3 = \mathbf{S}\underbrace{\mathbf{D}\mathbf{S}^{-1}\mathbf{S}}_{\text{cancels}}\underbrace{\mathbf{D}\mathbf{S}^{-1}\mathbf{S}}_{\text{cancels}}\mathbf{D}\mathbf{S}^{-1} = \mathbf{S}\mathbf{D}^3\mathbf{S}^{-1}.$$

More generally, for any integer $n \geq 0$,

$$\mathbf{A}^n = \mathbf{S}\mathbf{D}^n\mathbf{S}^{-1}.$$

Problem 16.1 Compute \mathbf{A}^{10} for $\mathbf{A} = \begin{pmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{pmatrix}$.

Solution : We have previously diagonalized this matrix:

$$\mathbf{A} = \mathbf{S}\mathbf{D}\mathbf{S}^{-1} \quad \text{where} \quad \mathbf{S} = \begin{pmatrix} 1 & -1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$$

$$\mathbf{D} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & -3 \end{pmatrix}.$$

This gives

$$\mathbf{A}^{10} = (\mathbf{S}\mathbf{D}\mathbf{S}^{-1})^{10}$$

$$= \mathbf{S} \mathbf{D}^{10} \mathbf{S}^{-1}$$

$$= \begin{pmatrix} 1 & -1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & (-3)^{10} & 0 \\ 0 & 0 & (-3)^{10} \end{pmatrix} \begin{pmatrix} 1 & -1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}^{-1}$$

16. Applications of diagonalization

[Hide Discussion](#)

Topic: Unit 2: Linear Algebra, Part 2 / 16. Applications of diagonalization

[Add a Post](#)

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

[Learn About Verified Certificates](#)

© All Rights Reserved