



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

43:44:37

< Previous

✓

✓

✓

✓

✓

✓

✓

✓

✓

✓

Next >

2. Vector addition

Bookmark this page

Problem Set A due Aug 18, 2021 20:30 IST Completed

2A-4

3.0/3 points (graded)

What is the vector \vec{v} that starts at the point $(1, 0)$ and ends at $(2, 2)$?

(Enter the vector in the form $[a, b]$. That is surround your vector by square brackets, and separate entries by a comma. Note that the entries of your vector must be numbers.)

$\vec{v} =$ ✓ Answer: [1,2]

Imagine the line segment from $(1, 0)$ to $(2, 2)$. Suppose we start at $(1, 0)$ and follow this line segment one third of the way to $(2, 2)$. What point do we end up at?

(Enter the ordered pair in round parentheses, e.g. (x, y) .)

✓ Answer: (4/3,2/3)

Imagine the line segment from $(1, 0)$ to $(2, 2)$. Suppose we start at $(1, 0)$ and go a distance of **0.5** along this line segment. What point do we end up at?

(Enter the ordered pair in parentheses, e.g. (x, y) .)

✓ Answer: (1+1/(2*sqrt(5)),1/sqrt(5))

? INPUT HELP

Solution:

The vector from a point (x_1, y_1) to a point (x_2, y_2) is given by subtracting the coordinates of the beginning point from the coordinates of the endpoint. In this case, we have

$$\vec{v} = \langle 2, 2 \rangle - \langle 1, 0 \rangle = \langle 1, 2 \rangle.$$

The vector \vec{v} above is the vector from $(1, 0)$ to $(2, 2)$. So to go a third of the way from $(1, 0)$ to $(2, 2)$, we should add $\frac{1}{3}\vec{v}$ to $\langle 1, 0 \rangle$. So the point we end up at is

$$(1, 0) + \left\langle \frac{1}{3}, \frac{2}{3} \right\rangle = \left\langle \frac{4}{3}, \frac{2}{3} \right\rangle.$$

To go a distance of **0.5** along the line segment from $(1, 0)$ to $(2, 2)$, we add a vector \vec{w} to $\langle 1, 0 \rangle$ where \vec{w} is in the same direction as \vec{v} and has length 0.5. The unit vector in the direction of \vec{v} is $\langle 1/\sqrt{5}, 2/\sqrt{5} \rangle$. So \vec{w} is given by

$$\vec{w} = \frac{1}{2} \langle 1/\sqrt{5}, 2/\sqrt{5} \rangle = \left\langle \frac{1}{2\sqrt{5}}, \frac{1}{\sqrt{5}} \right\rangle.$$

Adding this to $\langle 1, 0 \rangle$ means we end up at the point

$$(1,0) + \left(\frac{1}{2\sqrt{5}}, \frac{1}{\sqrt{5}}\right) = \left(1 + \frac{1}{2\sqrt{5}}, \frac{1}{\sqrt{5}}\right).$$

Submit

You have used 2 of 5 attempts

i Answers are displayed within the problem

2. Vector addition

Hide Discussion

Topic: Unit 2: Geometry of Derivatives / 2. Vector addition

Add a Post

Show all posts	▼	by recent activity	▼
?	<u>Mismatch in grading</u>	8	▼
	For 2A-4 I got the second answer correct but the grader marked as incorrect. I am not sure what I am missing here.		
💬	<u>error</u>	3	▼
	hey my answer was right but it wasnt accepted.		
?	<u>Answer check on part III of 2A-4</u>	6	▼
	Without giving away, I thought I had part III down pretty much in my head. If staff can see my answer, what am I missing?		
?	<u>Staff: Answer Check</u>	2	▼
	Hello, I was wondering if someone from the staff could check my answer to part 3; the problem is probably with me and not the gra...		
💬	<u>[Staff] Missing Backslash</u>	2	▼
💬	<u>The problem disappeared.</u>	3	▼
	I got an error message stating the following: "Could not format HTML for problem. Contact course staff in the discussion forum for a...		
?	<u>I had a * on an answer</u>	3	▼
	Hi Jennifer. I got an "*" on an answer. I cannot figure why. Can you have a look? Thanks		

< Previous

Next >



Calculator

Hide Notes



- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)