

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 3 Methods of Estimation](#) > [Lecture 9: Introduction to Maximum Likelihood Estimation](#) > [2. Review and Likelihood of a Gaussian Distribution](#)

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

2. Review and Likelihood of a Gaussian Distribution

Concept Check: Likelihoods of a Bernoulli, a Poisson, and a Gaussian Distribution



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Is the Likelihood Discrete or Continuous?

2/2 points (graded)

Setup:

Consider a **discrete** statistical model $M_1 = (\mathbb{Z}, \{\mathbf{P}_\theta\}_{\theta \in \mathbb{R}})$ and a **continuous** statistical model $M_2 = (\mathbb{R}, \{Q_\theta\}_{\theta \in \mathbb{R}})$. Let p_θ denote the pmf of \mathbf{P}_θ , and let q_θ denote the pdf of Q_θ . Assume that p_θ and q_θ both vary continuously with the parameter θ .

Let x_1, \dots, x_n be fixed natural numbers and y_1, \dots, y_n be fixed real numbers. Let $(L_1)_n$ denote the likelihood of the discrete model M_1 , and let $(L_2)_n$ denote the likelihood of the continuous model M_2 . Keeping x_1, \dots, x_n and y_1, \dots, y_n fixed, let's think of $(L_1)_n(x_1, \dots, x_n, \theta)$ and $(L_2)_n(y_1, \dots, y_n, \theta)$ as functions of θ .

Question

Decide whether the following claims about $(L_1)_n$ and $(L_2)_n$ are true or false.

The map $\theta \mapsto (L_1)_n(x_1, \dots, x_n, \theta)$ is a continuous function of θ .

☒ True☐ False

The map $\theta \mapsto (L_2)_n(y_1, \dots, y_n, \theta)$ is a continuous function of θ .

☒ True☐ False**Solution:**

Observe that

$$(L_1)_n(x_1, \dots, x_n, \theta) = \prod_{i=1}^n p_\theta(x_i), \quad (L_2)_n(y_1, \dots, y_n, \theta) = \prod_{i=1}^n q_\theta(y_i).$$

We are given that p_θ and q_θ are both continuous function of the parameter $\theta \in \mathbb{R}$. Since products of continuous functions are continuous, this implies that the maps $\theta \mapsto (L_1)_n(x_1, \dots, x_n, \theta)$ and $\theta \mapsto (L_2)_n(y_1, \dots, y_n, \theta)$ are continuous functions of the parameter $\theta \in \mathbb{R}$.

Remark: It may be confusing that even the likelihood of a discrete statistical model can be continuous. However, considering the likelihood of a Bernoulli (derived in a previous question),

$$L(x_1, \dots, x_n, p) = \prod_{i=1}^n p^{x_i} (1-p)^{1-x_i} = p^{\sum_{i=1}^n x_i} (1-p)^{n - \sum_{i=1}^n x_i}.$$

we can clearly see that the above varies continuously as a function of the *parameter*. This is also true for a host of other discrete models (for example, the Poisson model).

You have used 1 of 1 attempt

i Answers are displayed within the problem

Quiz: Likelihood of a Gaussian Statistical Model

3/3 points (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} N(\mu^*, (\sigma^*)^2)$ for some unknown $\mu^* \in \mathbb{R}, (\sigma^*)^2 > 0$. You construct the associated statistical model $(\mathbb{R}, \{N(\mu, \sigma^2)\}_{(\mu, \sigma^2) \in \mathbb{R} \times (0, \infty)})$.

The likelihood of this model can be written

$$L_n(x_1, \dots, x_n, (\mu, \sigma^2)) = \frac{1}{(\sigma\sqrt{2\pi})^C} \exp\left(-\frac{1}{A} \sum_{i=1}^C B_i\right)$$

where A depends on σ , B_i depends on μ and x_i . Find A, B_i and C .

(Choose a B_i that has coefficient 1 for the highest degree term in x_i .)

(Type **sigma** for σ , **mu** for μ , and **x_i** for x_i .)

$A =$ ✓

$2 \cdot \sigma^2$

$B_i =$ ✓

$x_i^2 + \mu^2 - 2 \cdot x_i \cdot \mu$

$C =$ ✓

n

STANDARD NOTATION

Submit

You have used 1 of 3 attempts

✓ Correct (3/3 points)

Discussion

Hide Discussion

Topic: Unit 3 Methods of Estimation:Lecture 9: Introduction to Maximum Likelihood Estimation / 2. Review and Likelihood of a Gaussian Distribution

Add a Post

Show all posts ▼

by recent activity ▼

? Should we consider cases where there may be an indicator function based on the parameter?

5

The prof uses a distribution such as this in one of the next few lectures and it leads me to one answer I think but it doesn't seem to be in the spirit of the question.

💬 [STAFF] Possible Error in Solution for Gaussian question

4

I believe the second term in the solution, after "the," is a possibly confusing typo. ****edit****: to clarify in title that this refers to Gaussian question.

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

Learn About Verified Certificates

© All Rights Reserved