

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 3 Methods of Estimation](#) > [Lecture 8: Distance measures between distributions](#)

14. Likelihood of a Discrete
> Distribution

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

14. Likelihood of a Discrete Distribution

Preparation: Equivalent Expressions for the pmf of a Bernoulli Distribution

1/1 point (graded)

Which of the following function $f(x)$, when restricted to the domain $x \in \{0, 1\}$, is equal to the pmf f of the probability distribution $\text{Ber}(p)$? Assume that $p \in (0, 1)$. (Choose all that apply.) (Recall that if $X \sim \text{Ber}(p)$, then $p = P(X = 1)$.)

☒ $f(x) = \begin{cases} p & \text{if } x = 1 \\ 1 - p & \text{if } x = 0 \end{cases}$

☒ $f(x) = p^x(1 - p)^{1-x}$

☒ $f(x) = xp + (1 - x)(1 - p)$

☐ $f(x) = \begin{cases} 1 & \text{if } x = 1 \\ 0 & \text{if } x = 0 \end{cases}$



Solution:

We will explain in the order of the choices.

- $f(x) = \begin{cases} p & \text{if } x = 1 \\ 1 - p & \text{if } x = 0 \end{cases}$ is correct. A random variable $X \sim \text{Ber}(p)$, by definition, has sample space $\{0, 1\}$ and satisfies $P(X = 1) = p$ and $P(X = 0) = 1 - p$. The given function is just a restatement of that definition.
- $f(x) = p^x(1 - p)^{1-x}$ is correct. Note that $f(1) = p$ and $f(0) = 1 - p$, so this is the same as the function considered in the first choice.
- $f(x) = xp + (1 - x)(1 - p)$ is correct. It also satisfies $f(1) = p$ and $f(0) = 1 - p$, so f is the same as the function considered in the first choice.
- $f(x) = \begin{cases} 1 & \text{if } x = 1 \\ 0 & \text{if } x = 0 \end{cases}$ is incorrect. This is actually the probability mass function of $\text{Ber}(1)$, but we have assumed $p \in (0, 1)$.

Submit

You have used 2 of 2 attempts

i Answers are displayed within the problem

Review: Statistical Model for a Bernoulli Distribution

3/3 points (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Ber}(p^*)$ for some unknown $p^* \in (0, 1)$. Let $(E, \{\text{Ber}(p)\}_{p \in \Theta})$ denote the corresponding statistical model. What is the smallest possible set that could be E ?

☐ $\{0\}$
☐ $\{-1, 1\}$
☒ $\{0, 1\}$

☐ \mathbb{R} 

The parameter space Θ can be written as an interval $[a, b]$. What is the smallest possible interval so that $\{\text{Ber}(p)\}_{p \in \Theta}$ represents all possible Bernoulli distributions?

 $a =$

Answer: 0.0

 $b =$

Answer: 1.0

Solution:


Since a Bernoulli random variable is either 0 or 1, the smallest possible sample space is $\{0, 1\}$.

If $\Theta = [0, 1]$, then $\{\text{Ber}(p)\}_{p \in [0, 1]}$ is the set of all possible Bernoulli distributions, as desired.

You have used 1 of 2 attempts

Answers are displayed within the problem

Likelihood of a Discrete Distribution



▶ 6:45 / 6:45

▶ 1.50x 🔊 🗑️ 📄 🗨️

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Concept Check: Interpreting the Likelihood

1/1 point (graded)

Let $(E, \{P_\theta\}_{\theta \in \Theta})$ denote a discrete statistical model. Let p_θ denote the pmf of P_θ . Let $X_1, \dots, X_n \stackrel{iid}{\sim} P_{\theta^*}$ where the parameter θ^* is unknown. Then the **likelihood** is the function

$$L_n : E^n \times \Theta \rightarrow \mathbb{R}$$

$$(x_1, \dots, x_n, \theta) \mapsto \prod_{i=1}^n p_\theta(x_i).$$

For our purposes, we think of x_1, \dots, x_n as observations of the random variables X_1, \dots, X_n .

Which of the following are true about the likelihood L_n ? (Choose all that apply.)

☒ It is the joint pmf of n iid samples from the distribution P_θ .

☒ It is a function of the sample $X_1 = x_1, \dots, X_n = x_n$.

☒ It is a function of the parameter θ , where θ ranges over all possible values of in the parameter space Θ .

☐ It is the joint pmf of n iid samples from the true distribution P_{θ^*} .



Solution:

We examine the choices in order.

- "It is the joint pmf of n iid samples from the distribution P_θ ." is correct. If $Y_1, \dots, Y_n \stackrel{iid}{\sim} P_\theta$, then by independence, the joint pmf of these variables is given by a product:

$$P(Y_1 = x_1, \dots, Y_n = x_n) = \prod_{i=1}^n p_\theta(x_i).$$

Remark 1: We use Y_i to denote these variables to differentiate from the samples X_i that come from the true distribution P_{θ^*} .

- "It is a function of the sample $X_1 = x_1, \dots, X_n = x_n$." is correct. To construct the likelihood, we observe samples $X_1 = x_1, \dots, X_n = x_n$ and then compute $L_n(x_1, \dots, x_n, \theta)$.
- "It is a function of the parameter θ , where θ ranges over all possible values of in the parameter space Θ " is correct. As θ varies over Θ , the likelihood $L_n(x_1, \dots, x_n, \theta)$ takes on different values. This is evident from the dependence on θ in the definition of the likelihood.

Remark 2: Later on we will maximize L_n (as a function of θ) to define the **maximum likelihood estimator**. Hence, it is a crucial property that the likelihood is a function of the parameter.

- "It is the joint pmf of n iid samples from the distribution P_{θ^*} ." is incorrect. The likelihood takes as input all possible θ , not just the true parameter θ^* . Note how the likelihood is defined for general θ , not just the true parameter θ^* .

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Likelihood of a Bernoulli Statistical Model

1/1 point (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Ber}(p^*)$ for some unknown $p^* \in (0, 1)$. Let $(E, \{\text{Ber}(p)\}_{p \in \Theta})$ denote the corresponding statistical model constructed in the previous question.

What is the likelihood L_n of this statistical model? (Choose all that apply.)

Hint: Use the pmf's in the second and third choices from the first problem on this page: "Preparation Equivalent Expressions for the pmf of a Bernoulli Distribution".

☒ $L_n(x_1, \dots, x_n, p) = p^{\sum_{i=1}^n x_i} (1-p)^{n - \sum_{i=1}^n x_i}.$

☐ $L_n(x_1, \dots, x_n, p) = p^{n - \sum_{i=1}^n x_i} (1 - p)^{\sum_{i=1}^n x_i}.$

☐ $L_n(x_1, \dots, x_n, p) = p^{\sum_{i=1}^n x_i} (1 - p)^{\sum_{i=1}^n x_i}.$

☒ $L_n(x_1, \dots, x_n, p) = \prod_{i=1}^n (x_i p + (1 - x_i)(1 - p))$



Solution:

We examine the choices in order.

- As shown in the previous problem, we can write the pmf of a Bernoulli as $x \mapsto p^x(1 - p)^{1-x}$. Hence,

$$L_n(x_1, \dots, x_n, p) = \prod_{i=1}^n p^{x_i} (1 - p)^{1-x_i} = p^{\sum_{i=1}^n x_i} (1 - p)^{n - \sum_{i=1}^n x_i}.$$

$$\begin{aligned} L(x_1, \dots, x_n, p) &= \prod_{i=1}^n p^{x_i} (1 - p)^{1-x_i} \\ &= p^{\sum_{i=1}^n x_i} (1 - p)^{n - \sum_{i=1}^n x_i}. \end{aligned}$$

Hence the first answer choice is correct.

- The second and third choices $L_n(x_1, \dots, x_n, p) = p^{n - \sum_{i=1}^n x_i} (1 - p)^{\sum_{i=1}^n x_i}$ and $L_n(x_1, \dots, x_n, p) = p^{\sum_{i=1}^n x_i} (1 - p)^{\sum_{i=1}^n x_i}$ are incorrect. Note that they are slight algebraic modifications of the first choice, so these formulas cannot be correct.
- If we use the expression $f(x) = xp + (1 - x)(1 - p)$ for the pmf of $\text{Ber}(p)$, then

$$L(x_1, \dots, x_n, p) = \prod_{i=1}^n (x_i p + (1 - x_i)(1 - p))$$

is, by definition, the likelihood. Hence, the last answer choice is also correct.

Remark: Although the last answer choice is formally correct, the formula is much more difficult to work with. It is often more convenient to use $p^{\sum_{i=1}^n x_i} (1 - p)^{n - \sum_{i=1}^n x_i}$ for the likelihood of a Bernoulli statistical model.

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 3 Methods of Estimation:Lecture 8: Distance measures between distributions / 14. Likelihood of a Discrete Distribution

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

Learn About Verified Certificates

© All Rights Reserved