



You are taking "5.2 Exam 2" as a timed exam. [Show more](#)

End My Exam

05:48:17



< Previous







Next >

3.7.1 Finger Exercise: Optimization
fundamentals and gradient descent

 Bookmark this page

Finger Exercises 4 due Sep 7, 2023 05:00 IST Completed

MO2.11

MO2.12

Consider the function

$$f(x) = x^4 + x^3 - 2x^2$$

(3.32)

over the interval $-2 \leq x \leq 1$. For your own reference, we suggest plotting this function on a grid with sufficiently fine spacing in x . From your plot, answer the following questions.

© All Rights Reserved



Problem: Determine number of mins and



4.0/4.0 points (graded)

[About](#)
How many minimums (local and global) does the function have on this interval? Don't count the endpoints.
[Affiliates](#)
[edX for Business](#)

[Open edX](#)

[Careers](#)
[News](#)

2

✓ Answer: 2

How many maximums (local and global) does the function have on this interval? Don't count the endpoints.

[Terms of Service & Honor Code](#)

[Privacy Policy](#)

[Accessibility Policy](#)

[Trademark Policy](#)

[Sitemap](#)

[Cookie Policy](#)

[Your Privacy Choices](#)

1

✓ Answer: 1



Answers are displayed within the problem

[Idea Hub](#)

[Contact Us](#)

[Help Center](#)

[Security](#)

[Media Kit](#)

2.0/2.0 points (graded)

Suppose the gradient descent method is applied

starting from the point $x = -0.1$. Assume that the step size is chosen small enough so that the method does converge to a minimum. What will the value of f



☐ $f \approx 0.0$

© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)

☐ $f \approx -0.4$

☒ $f \approx -0.8$

Discussions

Posting in discuss

All posts sorted by recent activity

[← Previous](#)

Next >