




< Previous













Next >

8.3.2 Skydiving

 Bookmark this page

MO2.4

For the skydiving example in Section 8.2.2, dividing Equation (8.8) by m gives

$$\frac{dV}{dt} = g - \frac{1}{2} \frac{\rho_a V^2 A_{\text{ref}} C_D}{m}$$

(8.38)

which in terms of the general IVP form gives a scalar ($M = 1$) system of equations with

$$u = V \quad f = g - \frac{1}{2} \frac{\rho_a u^2 A_{\text{ref}} C_D}{m}$$

(8.39)

< Previous

Next >

Discussions

All posts sorted by recent activity



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)