EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the Privacy Policy.





MITx: 6.86x

Course > Generalizations (2 weeks)

Machine Learning with Python-From Linear Models to Deep Learning

<u>Help</u>



<u>sandipan_dey</u>

Unit 1 Linear Classifiers and

Lecture 3 Hinge loss, Margin


> boundaries and Regularization

> 2. Introduction

2. Introduction Introduction



Learning as optimization



for selecting the parameters theta and theta naught.

That's a balance between the loss, how examples

fit within this ideal notion, and regularization, our preference towards large mounting solutions.

So we will find--

we will formalize-- the objective function and then find parameters theta and theta

nought that optimize, minimize, this objective function.

6:19 / 6:19

▶ Speed 1.50x





End of transcript Clin to the start

Video Download video file **Transcripts** Download SubRip (,srt) file Download Text (.txt) file


Review: Distance from a Line to a Point

1 point possible (graded) Consider a line L in \mathbb{R}^2 given by the equation

$$L: \theta \cdot x + \theta_0 = 0$$

where θ is a vector normal to the line L. Let the point P be the endpoint of a vector x_0 (so the coordinates of P equal the components of x_0).

What is the the shortest distance d between the line L and the point P? Express d in terms of θ, θ_0, x, x_0 .



$$d =$$

$$\frac{|\theta \cdot x + \theta_0|}{||\theta||}$$

$$egin{array}{ccc} & rac{| heta \cdot x_0 + heta_0|}{|| heta||}
ightharpoonup \ \end{array}$$

$$\bigcirc \frac{|\theta \cdot \theta_0 + \theta_0|}{||\theta||}$$

$$ullet | heta \cdot x_0 + heta_0|$$

Solution:

If there is no offset θ_0 , The distance d is the projection from x_0 to θ , which is $\frac{|x_0\cdot\theta|}{||\theta||}$ (definition of projection). With the offset θ_0 added, d is $\frac{|x_0\cdot\theta+\theta_0|}{||\theta||}$. Thus the distance from a $L:\theta\cdot x+\theta_0=0$ to the point $P=x_0$ is given by $\frac{|\theta\cdot x_0+\theta_0|}{||\theta||}$.

Submit

You have used 0 of 3 attempts

1 Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 1 Linear Classifiers and Generalizations (2 weeks):Lecture 3 Hinge loss, Margin boundaries and Regularization / 2. Introduction

Add a Post

Sho	by recent activ	vity ▼
2	Can someone explain the distance problem please? I don't understand the answer to the distance question. Can someone please explain, or provide references where I can learn more about the equations and meaning of the s	11
?	Reset grader In all fairness, I request staff to reset grader for unit 1 lecture 3 and Homework 1 since the timeline is extended by a day. Thanking you.	6
∀	What is the significance of the "regularization term" in context of this "optimization problem"? Lesson says it's the favoring of margin boundaries that are far apart from the decision boundary. I am not sure about two things here - 1. What having margin boundaries tha	2
∀	What exactly is an "optimization problem" and why do we need it here? I don't think the lesson clarifies this. I am confused why we are even defining all of this.	2
∀	Clarification regarding the loss function at 5:14, Professor says "we might start violating the preference that all the training examples are outside this fat boundary. They may go within the fat boundary or even be m	5
2	Staff - Caption [INAUDIBLE] Please correct the caption [INAUDIBLE], because my studies are only with the legend and in some lectures they are with this message.	4
∀	Course Progress Does Not Give Credit for Correct Answer to "Review: Distance from a Line to a Point" Question Grader gives me 12 points for Excercise 3, but Course Progress gives me only 11 points! After L get correct answer to "Review: Distance from a Line to a Point" question, I see L	2

∀	Distance from a Line to a Point - always abs(nominator)? Can this value be negative? Or it's always expressed as abs?	3
∀	Equidistant Margin Boundary Is it necessary to have equidistant margin boundary? If yes, then what advantage it will offer compare to non- equidistant margin boundary?	3
∀	the coordinates of P equal the components of x0? If x0 is a vector and P is its end point, then P doesn't equal coordinates (components) of the vector unless the beginning of the vector is in origin. Can you please clarify? I got t	2
∀	Latex typo in the explanation	2
	Learn About Verified Certificates	

© All Rights Reserved