



[Course](#) > [Unit 1: Fourier Series](#) > [Part A Homework 1](#) > 1. Lecture 1

**Audit Access Expires Jun 24, 2020**

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 1. Lecture 1

The following can be done after Lecture 1.

1-1

5/5 points (graded)

Recall that a function  $f(t)$  is called periodic of period  $P$  if  $f(t + P) = f(t)$  for all  $t$ . True or false: The function  $\cos 5t$  is periodic of period  $2\pi$ .

☒ True

☐ False



**Solution:**

True, because



$$\cos(5(t + 2\pi)) = \cos(5t + 10\pi) = \cos 5t,$$

since  $10\pi$  is an integer times  $2\pi$ . (The **smallest** period of  $\cos 5t$  is  $2\pi/5$ , but that is not what this problem is asking about.)

Submit

You have used 1 of 1 attempt

 Answers are displayed within the problem

1-2

5/5 points (graded)

The function  $\sin^2 t$  is

☒ even but not odd.

☐ odd but not even.

☐ both even and odd.

☐ neither even nor odd.



**Solution:**

Even but not odd. It is even because  $\sin^2(-t) = (-\sin t)^2 = \sin^2 t$ . It is not odd, since  $\sin^2(-t) = -\sin^2 t$  is not true for **all**  $t$  (for example, it fails when  $t = \pi/2$ ).

Submit

You have used 1 of 2 attempts



**i** Answers are displayed within the problem

1-3

10/10 points (graded)

Let  $f(t)$  be the function of period  $2\pi$  such that

$$f(t) = \begin{cases} 0 & \text{for } -\pi < t \leq 0; \\ 3 & \text{for } 0 < t \leq \pi \end{cases}.$$

What is the value of the constant coefficient in the Fourier series of  $f$ ?

3/2

✓ Answer: 3/2

$\frac{3}{2}$

**Solution:**

The constant term is  $\frac{a_0}{2}$ , where

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) dt = \frac{1}{\pi} \left( \int_{-\pi}^0 0 dt + \int_0^{\pi} 3 dt \right) = \frac{1}{\pi} \int_0^{\pi} 3 dt = 3.$$

Therefore the constant term is  $\frac{a_0}{2} = \frac{3}{2}$ .

Alternative solution: The answer is the average value of  $f(t)$  on one period, which is  $3/2$ , the average of 0 and 3, because the interval on which  $f(t) = 0$  has the same length as the interval on which  $f(t) = 3$ .

Submit

You have used 1 of 10 attempts



**i** Answers are displayed within the problem

1-4

5/5 points (graded)

Let  $f(t)$  be the periodic function of period  $2\pi$  such that  $f(t) = |t|$  for  $-\pi \leq t < \pi$ . What is the coefficient of  $\sin 7t$  in the Fourier series for  $f$ ?

**Note: You must use a star to denote multiplication; e.g.  $7 * x = 7x$ . Use  $\wedge$  to denote exponentiation; e.g.  $e \wedge x = e^x$ . A slash denotes division; e.g.  $1/2 = 0.5$ . Please type pi rather than a numerical approximation for  $\pi$ .**

0

✓ Answer: 0

0

**Solution:**

The answer is 0, because  $f(t)$  is an even function.

Submit

You have used 1 of 10 attempts

**i** Answers are displayed within the problem

## 1. Lecture 1

Hide Discussion

**Topic:** Unit 1: Fourier Series / 1. Lecture 1

Add a Post

Show all posts ▼

by recent activity ▼

✓ [staff] terminology in 1-3

Since the 'a' coefficients are a sub n, isn't the constant coefficient a sub 0?

2

Learn About Verified Certificates

