



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

44:25:12



< Previous



Next >

5. Practice

Bookmark this page



Calculator



Hide Notes

Problem Set B due Aug 4, 2021 20:30 IST Completed



Practice


A second example, level curves

1/1 point (graded)


What are the level curves of the function $\sin(x + t)$?

Use the convention that the x axis is the vertical axis, and the t axis is the horizontal axis.


☐




☒



☐





☐



Calculator

Hide Notes



Solution:

To understand the level curves, it is helpful to look at specific levels and understand what is happening.

First let us consider the level curve $\sin(x+t) = 0$ Note that this happens when $x+t = \pi k$ where k is any integer. Therefore the level curves for $\sin(x+t) = 0$ is a collection of lines

$$\{t = \pi k - x | k \text{ an integer}\}$$

Note that these lines are all parallel to each other, with intercepts along the t axis differing by integer values of π .

We see that the level curves are very similar to the level curves for $\sin(x-t)$, except that the level curves are now lines parallel to $t = -x$ instead of $t = x$.

Submit

You have used 1 of 3 attempts

Answers are displayed within the problem

A second example, time perspective

1/1 point (graded)

What best describes how we observe the function $\sin(x+t)$ as a function that changes over time?

- ☐ A sine function that appears to travel to the right over time.
- ☒ A sine function that appears to travel to the left over time.
- ☐ A sine function whose frequency increases over time.
- ☐ A sine function whose frequency decreases over time.

☐ A sine function whose amplitude increases over time.

☐ A since function whose amplitude decreases over time.



Solution:

This is similar to the first example, except at every time t_0 , the sine function is shifted to the left by t_0 .

Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

A third example

1/1 point (graded)

What best describes how we observe the function $e^{-t} \sin(x)$ as a function that changes over time?

☐ A sine function that appears to travel to the right over time.

☐ A sine function that appears to travel to the left over time.

☐ A sine function whose frequency increases over time.

☐ A sine function whose frequency decreases over time.

☐ A sine function whose amplitude increases over time.


☒ A sine function whose amplitude decreases over time.



Solution:

Note that at $t = 0$, we get the function $\sin(x)$. At a later time t , the function is $e^{-t} \sin(x)$. Therefore the amplitude of the sine function is being multiplied by a number $e^{-t} < 1$ for all $t > 0$. The larger t is, the smaller e^{-t} becomes. Therefore as t increases, the amplitude of the sine function is decreasing.

Below you see the function $\sin(x)$ in blue, and the function $e^{-t} \sin(x)$ animated over time in orange.



Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

5 Practice

< Previous

Next >

© All Rights Reserved



edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers
- News

Legal

- Terms of Service & Honor Code
- Privacy Policy
- Accessibility Policy
- Trademark Policy
- Sitemap

Connect

- Blog
- Contact Us
- Help Center
- Media Kit
- Donate



