



Bookmarks

- ▶ [Module 1: The Basics of R and Introduction to the Course](#)
- ▶ [Entrance Survey](#)
- ▶ [Module 2: Fundamentals of Probability, Random Variables, Distributions, and Joint Distributions](#)
- ▶ [Module 3: Gathering and Collecting Data, Ethics, and Kernel Density Estimates](#)
- ▶ [Module 4: Joint, Marginal, and Conditional Distributions & Functions of Random Variable](#)

Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation > The Sample Mean, Central Limit Theorem, and Estimation > More on Criteria for Assessing Estimators - Quiz

More on Criteria for Assessing Estimators - Quiz

🔖 Bookmark this page

Question 1

1/1 point (graded)

True or False: To prove an estimator is unbiased, you need to know the value of the parameter it is trying to estimate.

☐ a. True

☒ b. False ✓

Explanation

You can prove an estimator to be unbiased through a mathematical proof, without knowing the underlying value of the parameter. In Segment 4 of this lecture, Professor Ellison mathematically proved that the sample mean is an unbiased estimate of the mean.


Submit

You have used 1 of 1 attempts


► [Module 5: Moments of a Random Variable, Applications to Auctions, & Intro to Regression](#)

▼ [Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation](#)


[Human Subjects and Special Distributions](#)

Finger Exercises due Nov 07, 2016 at 05:00 IST 

[The Sample Mean, Central Limit Theorem, and Estimation](#)

Finger Exercises due Nov 07, 2016 at 05:00 IST 

[Module 6: Homework](#)

Homework due Oct 31, 2016 at 05:00 IST 

► [Exit Survey](#)

✓ Correct (1/1 point)

Question 2

1 point possible (graded)

For the example distribution in lecture, which of the following methods would result in an unbiased estimate of θ ?

☐ a. Compute the median of the sample and multiply by 2 ✓

☒ b. Compute the maximum (nth order statistic) of the sample

☐ c. Have R generate a random value from the underlying distribution.

☒ d. Compute the sample mean and multiply by 2 ✓



Explanation

The expected value of the sample mean and the sample median will be $\frac{\theta}{2}$. Multiplying that by 2 gives us an expected value of θ . Having R generate a random value from the underlying distribution would have an expected value of only $\frac{\theta}{2}$. Computing the maximum (n^{th} order statistic of the sample) would have an expected value of less than θ . The n^{th} order statistic will always be less than or equal to θ , so it will be a downwardly biased estimate of θ .

[Submit](#)

You have used 2 of 2 attempts

✖ Incorrect (0/1 point)

Discussion

Topic: Module 6 / More on Criteria for Assessing Estimators - Quiz

[Show Discussion](#)

© All Rights Reserved



© 2016 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY
OPENedX



