



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

44:53:54



< Previous



## 14. Partial derivatives for hikers

🔖 Bookmark this page



Calculator



Hide Notes

Lecture due Aug 4, 2021 20:30 IST Completed



## Synthesize

Now, we'll see how what we've learned so far connects to the problem involving Hikers 1 and 2.

## Hikers computation 1

2.0/2 points (graded)

Let  $f(x, y) = x^2 + y^2$ . Compute:

$f_x(x, y) =$

2\*x

✓ Answer: 2\*x

$f_y(x, y) =$

2\*y

✓ Answer: 2\*y

? INPUT HELP

## Solution:

To compute  $f_x(x, y)$ , we hold  $y$  constant and differentiate with respect to  $x$ :

$$f_x(x, y) = \frac{\partial}{\partial x}(x^2 + y^2) = 2x.$$

Notice that since  $y$  is treated as a constant, the derivative with respect to  $x$  of  $y^2$  is 0.

To compute  $f_y(x, y)$ , we hold  $x$  constant and differentiate with respect to  $y$ :

$$f_y(x, y) = \frac{\partial}{\partial y}(x^2 + y^2) = 2y.$$

Notice that since  $x$  is treated as a constant, the derivative with respect to  $y$  of  $x^2$  is 0.

Submit

You have used 1 of 5 attempts

❗ Answers are displayed within the problem

## Hikers computation 2

2/2 points (graded)

Let  $f(x, y) = x^2 + y^2$ . Compute:

$f_x(-1, 1) =$

-2

✓ Answer: -2

$f_x(-2, 0) =$

-4

✓ Answer: -4

## Solution:

From the previous problem, we have

Calculator

Hide Notes

$$f_x(x,y) = 2x.$$

Then


$$f_x(-1,1) = 2(-1) = -2$$

and

$$f_x(-2,0) = 2(-2) = -4.$$

Submit

You have used 1 of 5 attempts

 Answers are displayed within the problem

Recall that, when Hiker 1 started at  $(-1,1)$  and moved in the positive  $x$ -direction, we deduced from the level curves that the hiker was traveling downhill.

Hiker 2 started at  $(-2,0)$  and moved in the positive  $x$ -direction. Recall that we deduced from the level curves that Hiker 2 was moving downhill more steeply than Hiker 1.












Think about the sign and magnitude of your answers above. How do these correspond to your answers about Hiker 1 and Hiker 2? Use the discussion forum to explain.




14. Partial derivatives for hikers

Hide Discussion

Topic: Unit 1: Functions of two variables / 14. Partial derivatives for hikers

Add a Post

Show all posts	by recent activity
<div><div></div><div><a href="#">Partial derivatives for hikers</a></div><div>The partial derivative represents the slope of the tangent line that is parallel to one of the coordinate axes. Since the slope is negativ...</div><div>1</div></div>	
<div><div></div><div><a href="#">Hiker 1 vs Hiker 2</a></div><div>Good morning, I think I understood something. With the partial derivatives something similar happens to what happens with the deri...</div><div>2</div></div>	
<div><div></div><div><a href="#">Slope behavior</a></div><div>Since the partial derivative shows us how *f* changes, the negative sign tells us that they are going downhill, while the number how...</div><div>2</div></div>	
<div><div></div><div><a href="#">Steepness Magnitude</a></div><div>From the deduction, Hiker 1 which was moving down-hill had a lower negative value which will confirm that it's less steeper than Hik...</div><div>1</div></div>	
<div><div></div><div><a href="#">Steepness</a></div><div></div><div>1</div></div>	
<div><div></div><div><a href="#">Partial rate of change?</a></div><div>The partial derivative gives us the instantaneous rate of change when moving in one axis. In this case it is larger in the negative dire...</div><div>1</div></div>	
<div><div></div><div><a href="#">Warning about mobile version of page</a></div><div></div><div>3</div></div>	
<div><div></div><div><a href="#">An instantaneous 'rate of change'</a></div><div></div><div>1</div></div>	
<div><div></div><div><a href="#">Partial derivatives for hikers</a></div><div>The partial derivatives at (-2,0) and (-1,0) are negative, suggesting they are moving downhill. The absolute value of Hiker 1's derivati...</div><div>1</div></div>	
<div><div></div><div><a href="#">Mountain Analogy</a></div><div></div><div>5</div></div>	
<div><div></div><div><a href="#">Rate of change</a></div><div></div><div></div></div>	

	<u>Relationship of Partial Derivatives to the Contour Plot</u> The signs of the partial derivativs at (-2,0) and (-1,0) are negative, indicating that they are descending. The differences in the absolu...	1
	<u>more negative answer</u> Since hiker 2 has a MORE negative answer means that the value of z (i.e. altitude) is decreasing faster when moving in the x direction.	2
	<u>rate of descent</u> The walker at (-2,0) experiences a larger rate of descent for a given change in the x-direction than the walker at (-1,1)	1



## edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

## Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

## Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)





© 2021 edX Inc. All rights reserved.  
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)