

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).




[Course](#) > [Unit 3 Methods of Estimation](#) > [Likelihood Estimation](#) > [Lecture 9: Introduction to Maximum Likelihood Estimation](#) > 11. Strictly Concave Functions and Unique Maximizer

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

## 11. Strictly Concave Functions and Unique Maximizer

### Strictly Concave Functions and Unique Maximizer



▶ 4:47 / 4:47

▶ 1.50x 🔊 🗖 📄 🗑

### Video

[Download video file](#)

### Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

## Concavity and Convexity in Higher Dimensions II

2/2 points (graded)

As in the problem on the previous page,  $f(x, y) = -2x^2 + \sqrt{2}xy - \frac{5}{2}y^2$ . Based on your answer to the question, which of the following is true?

☐  $f$  has a unique (global) minimizer.

☒  $f$  has a unique (global) maximizer.

☐  $f$  has more than one (global) minimizer

☐  $f$  has more than one (global) maximizer



Where is the critical point of  $f$ ? (If there is more than one critical point, just enter one of them.)

(Enter your answer as a vector, e.g., type **[3,2]** for the vector  $\begin{pmatrix} 3 \\ 2 \end{pmatrix}$ , which corresponds to the point  $(3, 2)$  on the  $(x, y)$ -plane).

Critical point of  $f$ :

✓ Answer: [0,0]

STANDARD NOTATION

### Solution:

Since  $f$  is twice-differentiable and strictly concave, we know there will be a unique global maximum.

The critical points of  $f$  are those points where  $\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}\right) = 0$ . We compute that

$$\nabla f = (-4x + \sqrt{2}y, -5y + \sqrt{2}x),$$

which is 0 if and only if  $x = y = 0$ .

Submit

You have used 1 of 2 attempts

**i** Answers are displayed within the problem

## Concavity Concept Check

0/1 point (graded)

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice-differentiable function such that the Hessian matrix  $\mathbf{H}f(0,0)_{1,1} > 0$ . Is  $f$  concave?

☐ Yes

☒ No ✓

☐ Not possible to determine from given information

✗

### Solution:

Recall that  $f$  is concave at  $(0,0)$  if for all  $(x,y) \in \mathbb{R}^2$ ,

$$(x,y) \mathbf{H}f(0,0) \begin{pmatrix} x \\ y \end{pmatrix} < 0.$$

By expanding and using the definition of the Hessian, we see that

$$(x, y) \mathbf{H} f(0, 0) \begin{pmatrix} x \\ y \end{pmatrix} = x^2 \frac{\partial^2}{\partial x^2} f(0, 0) + xy \left( \frac{\partial^2}{\partial x \partial y} f(0, 0) + \frac{\partial^2}{\partial y \partial x} f(0, 0) \right) + y^2 \frac{\partial^2}{\partial y^2} f(0, 0).$$

By assumption, we know that  $\frac{\partial^2}{\partial x^2} f(0, 0) > 0$ . Hence,

$$(1, 0)^T \mathbf{H} f(0, 0) \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \frac{\partial^2}{\partial x^2} f(0, 0) > 0.$$

This violates the definition of concavity, so the correct response is "No."

Submit

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

**Note:** The following problem will be presented in lecture, but we encourage you to attempt it first.

## Intuition for Optimizing Concave Functions

1/1 point (graded)

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a twice-differentiable function that has a critical point and is strictly concave. Recall that the critical point is a **unique maximizer** of  $f$ .

You choose an initial guess for the maximizer  $x_0 = 0$  (which may be very far from the true maximizer). You compute the derivative and observe that  $f'(x_0) < 0$ . Where is the maximizer of  $f$ ?

☒ To the left of  $x_0 = 0$

☐ To the right of  $x_0 = 0$ ☐ Very far from  $x_0$ ☐ Very close to  $x_0$ **Solution:**

Graphically, a strictly concave function that has a critical point looks like a hill. If you are to the right of the peak (*i.e.*, the maximum), then the hill is sloping downward. If you are to the left of the peak, then the hill is sloping upward.

More formally, a differentiable function that is strictly concave has a strictly decreasing derivative. The first derivative is zero at the critical point, so it must be positive to the left of the maximum (which implies the function is increasing) and negative to the right of the maximum (which implies the function is decreasing).

Thus the first choice, "To the left of  $x_0 = 0$ ", is correct.

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Discussion

**Topic:** Unit 3 Methods of Estimation:Lecture 9: Introduction to Maximum Likelihood Estimation / 11. Strictly Concave Functions and Unique Maximizer

Add a Post

Show all posts ▼

by recent activity ▼

[What does  \$H\(f\(0,0\)\)\_{1,1}\$  mean?](#)

3

**Audit Access Expires Dec 24, 2019**

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

[Learn About Verified Certificates](#)

© All Rights Reserved