

edX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#). ✕



[Unit 2 Nonlinear Classification,](#)  
[Linear regression, Collaborative](#)  
[Course](#) > [Filtering \(2 weeks\)](#)

1. Collaborative Filtering, Kernels,  
 > [Homework 3](#) > Linear Regression

## 1. Collaborative Filtering, Kernels, Linear Regression

In this question, we will use the alternating projections algorithm for low-rank matrix factorization, which aims to minimize

$$J(U, V) = \underbrace{\frac{1}{2} \sum_{(a,i) \in D} (Y_{ai} - [UV^T]_{ai})^2}_{\text{Squared Error}} + \underbrace{\frac{\lambda}{2} \sum_{a=1}^n \sum_{j=1}^k U_{aj}^2 + \frac{\lambda}{2} \sum_{i=1}^m \sum_{j=1}^k V_{ij}^2}_{\text{Regularization}}.$$

In the following, we will call the first term the squared error term, and the two terms with  $\lambda$  the regularization terms.

Let  $Y$  be defined as

$$Y = \begin{bmatrix} 5 & ? & 7 \\ ? & 2 & ? \\ 4 & ? & ? \\ ? & 3 & 6 \end{bmatrix}$$

$D$  is defined as the set of indices  $(a, i)$ , where  $Y_{a,i}$  is not missing. In this problem, we let  $k = \lambda = 1$ . Additionally,  $U$  and  $V$  are initialized as  $U^{(0)} = [6, 0, 3, 6]^T$ , and  $V^{(0)} = [4, 2, 1]^T$ .

### 1. (a)

1.0/1 point (graded)

Compute  $X$ , the matrix of predicted rankings  $UV^T$  given the initial values for  $U$  and  $V$ .

[[24, 12, 6], [0, 0, 0], [12, 6, 3], [24, 12, 6]]

✓ Answer: [[24, 12, 6], [0, 0, 0], [12, 6, 3], [24, 12, 6]]

#### Solution:

- the predicted rankings should be the matrix product between  $U$  and  $V^T$ .

$$X = UV^T = \begin{bmatrix} 24 & 12 & 6 \\ 0 & 0 & 0 \\ 12 & 6 & 3 \\ 24 & 12 & 6 \end{bmatrix}$$

You have used 2 of 3 attempts

**i** Answers are displayed within the problem

1. (b)

2/2 points (graded)

Compute the squared error term, and the regularization terms in for the current estimate  $X$ .Enter the squared error term (including the factor  $1/2$ ):

✓ Answer: 255.5

Enter the regularization term (the sum of all the regularization terms):

✓ Answer: 51

**Solution:**

$$\begin{aligned} J_{\text{square}} &= \sum_{i,j \in D} (Y_{ij} - X_{ij})^2 / 2 \\ &= \frac{1}{2} \left( (5 - 24)^2 + (7 - 6)^2 + (2 - 0)^2 + (4 - 12)^2 + (3 - 12)^2 + (6 - 6)^2 \right) = 255.5 \end{aligned}$$

$$\begin{aligned}
 J_{\text{reg}} &= \frac{\lambda}{2} \|U\|_F^2 + \frac{\lambda}{2} \|V\|_F^2 \\
 &= \frac{\lambda}{2} \sum_{a=1}^n (U_a)^2 + \frac{\lambda}{2} \sum_{i=1}^m (V_i)^2 = 51
 \end{aligned}$$

Submit

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

1. (c)

1.0/1 point (graded)

Suppose  $V$  is kept fixed. Run one step of the algorithm to find the new estimate  $U^{(1)}$ .

Enter the  $U^{(1)}$  as a list of numbers,  $[U_1^{(1)}, U_2^{(1)}, U_3^{(1)}, U_4^{(1)}]$ :

[1.5, 0.8, 0.94117647, 2.0]

✓ Answer: [3/2, 4/5, 16/17, 2]

**Solution:**

With  $V$  fixed as  $[4, 2, 1]^T$ , we can represent prediction  $X$  as:

$$X = UV^T = \begin{bmatrix} 4U_1 & 2U_1 & 1U_1 \\ 4U_2 & 2U_2 & 1U_2 \\ 4U_3 & 2U_3 & 1U_3 \\ 4U_4 & 2U_4 & 1U_4 \end{bmatrix}$$

Let  $D$  be the set of index of observation, the estimate  $U^{(1)}$  should be:

$$\begin{aligned} U^{(1)} &= \arg \min_U J(U) \\ &= \arg \min_U \sum_{(a,i) \in D} (Y_{ai} - (UV)_{ai})^2 / 2 + \sum_{a=1}^4 \frac{\lambda}{2} \|U_a\|^2 \\ &= \arg \min_U [(5 - 4U_1)^2 + (7 - U_1)^2 + (2 - 2U_2)^2 + (4 - 4U_3)^2 + (3 - 2U_4)^2 + (6 - U_4)^2] / 2 + \sum_{a=1}^4 \frac{1}{2} U_a^2 \end{aligned}$$

To minimize this loss, we take the gradient with respect to  $U$  and equate it to zero.

$$0 = \nabla J(U) = \begin{pmatrix} -4(5 - 4U_1) - (7 - U_1) + U_1 \\ -2(2 - 2U_2) + U_2 \\ -4(4 - 4U_3) + U_3 \\ -2(3 - 2U_4) - (6 - U_4) + U_4 \end{pmatrix} = \begin{pmatrix} -27 + 18U_1 \\ -4 + 5U_2 \\ -16 + 17U_3 \\ -12 + 6U_4 \end{pmatrix}$$

Hence,

$$U_1^{(1)} = \frac{3}{2}$$
$$U_2^{(1)} = \frac{4}{5}$$
$$U_3^{(1)} = \frac{16}{17}$$
$$U_4^{(1)} = 2$$

Submit

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Discussion


Hide Discussion

**Topic:** Unit 2 Nonlinear Classification, Linear regression, Collaborative Filtering (2 weeks):Homework 3 / 1. Collaborative Filtering, Kernels, Linear Regression

Add a Post


Show all posts ▼

by recent activity ▼

 I can see regulation loss here, but what's the symptom of over fitting for Collaborative Filtering?

9

 Community TA

 [Staff] 1(a) Incomplete formulation of question

12

<p>🗨 <u>Staff my first answer was on the 3 percent difference</u></p> <p><u>I am learning and that is what matters to me, but my sloppy handwriting made me see a ? as a 7 (seven) and then my answer differs. On other co...</u></p>	2
<p>🗨 <u>Sage way.</u></p> <p><u>If you are tired of computing using pen and paper... sage: var('u1,u2,u3,u4,v1,v2,v3')(u1, u2, u3, u4, v1, v2, v3) sage: U = matrix([u1,u2,u3,u4]).tran...</u></p> <p>👤 <u>Community TA</u></p>	23
<p>🗨 <u>Hints</u></p> <p><u>I'm very stuck on where to even begin on this problem. Can anyone give me a hint on where to start with 1a? It would be very much appreciated,...</u></p>	9
<p>🗨 <u>matrix produce typo</u></p> <p><u>On the answer</u></p>	2
<p>🗨 <u>[Staff] Clarification 1 (b)</u></p> <p><u>For the regularization term in 1. (b) it would be helpful to make clear if we should include the half factor or not. I was doing something wrong an...</u></p>	2
<p>🗨 <u>[STAFF] Problem 1. (a)</u></p> <p><u>Hi, Could you please take a look at my solution to the first problem ? I believe I got the correct figures but somehow my answer was marked 'Inco...</u></p>	7
<p>✅ <u>Why only one regularization term?</u></p> <p><u>Why do we only use one regularization for both the U and V matrices? Would it make sense to use unique regularization terms for each matrix?</u></p>	3
<p>✅ <u>[Hint] Deciding Which U0 V0 and Lambda to Choose</u></p> <p><u>how do we decide which U0,V0 and lambda to choose? We decide which rank to use on development set. I dont know what is development set b...</u></p>	2
<p>✅ <u>is it OK to write in (c) answer list elements as fractions a/b ?</u></p> <p><u>subj</u></p>	4
<p>✅ <u>Is there an elegant formula for the objective function using linear algebra notation?</u></p>	5
<p>✅ <u>Can someone from staff please check my answer for 1(c)?</u></p> <p><u>I checked my calculation thrice but always getting red light from the grader.</u></p>	2

