







< Previous	 	 			Next >
-------------------------------	---	---	---	---	---------------------------

9.5.3 Bases for Subspaces

 Bookmark this page

Week 9 due Dec 9, 2023 18:12 IST Completed

9.5.3 Bases for Subspaces

Video

Summary

We will see next that a basis is a spanning set with minimum number of vectors.

▶

these bases have to have the same number of vectors in them.

33 / 33

▶ 5:47 / 5:47

▶ 2.0x

◀

⌂

CC

“

that you have exhibited the entire subspace.

I guess that should be an n here.

Anyway, you get the idea.

What we're going to see next is that a basis is a spanning set with a minimum number of vectors, and indeed,

if you have two different basis for the same subspace then

these bases have to have the same number of vectors in them.

Video

 [Download video file](#)

Transcripts

 [Download SubRip \(.srt\) file](#)

 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)

Read Unit 9.5.3 of the notes. [\[LINK\]](#)

☒ Done



Submit

✓ Correct

Discussion

Topic: Week 9 / 9.5.3

Hide Discussion

Add a Post

Show all posts

by recent activity ▼

☒ Linearly independent matrix and invertible matrix

Hi, At 2:15 Dr. Robert van de Geijn said "we know that these vectors are linearly independent because otherwise matrix A would not be invertible".



 Calculator

? # of equations vs unknowns

2

At 3:13, professor said "if you have fewer equations than unknowns, then we can find a non-0 vector \mathbf{x} such that $\mathbf{V}\mathbf{x} = \mathbf{0}$ ". How did we prove...

Homework 9.5.3.1

1/1 point (graded)

The vectors $\{\mathbf{e}_0, \mathbf{e}_1, \dots, \mathbf{e}_{n-1}\} \subset \mathbb{R}^n$ are a basis for \mathbb{R}^n .

TRUE



Answer: TRUE

Clearly, $\text{Span}(\mathbf{e}_0, \mathbf{e}_1, \dots, \mathbf{e}_{n-1}) = \mathbb{R}^n$. Now, the identity $\mathbf{I} = \left(\begin{array}{c|c|c|c} \mathbf{e}_0 & \mathbf{e}_1 & \cdots & \mathbf{e}_{n-1} \end{array} \right)$. Clearly, $\mathbf{I}\mathbf{x} = \mathbf{0}$ only has the solution $\mathbf{x} = \mathbf{0}$. Hence the columns of \mathbf{I} are linearly independent which means the vectors $\{\mathbf{e}_0, \mathbf{e}_1, \dots, \mathbf{e}_{n-1}\}$ are linearly independent.

Submit

< Previous

Next >

© All Rights Reserved



edX

[About](#)

[Affiliates](#)

[edX for Business](#)

[Open edX](#)

[Careers](#)

[News](#)

Legal

[Terms of Service & Honor Code](#)

[Privacy Policy](#)

[Accessibility Policy](#)

[Trademark Policy](#)

[Sitemap](#)

[Cookie Policy](#)

[Your Privacy Choices](#)

Connect

[Idea Hub](#)

[Contact Us](#)

[Help Center](#)

[Security](#)

[Media Kit](#)



Calculator



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)