



Bookmarks

- ▶ [Module 1: The Basics of R and Introduction to the Course](#)
- ▶ [Entrance Survey](#)
- ▶ [Module 2: Fundamentals of Probability, Random Variables, Distributions, and Joint Distributions](#)
- ▶ [Module 3: Gathering and Collecting Data, Ethics, and Kernel Density Estimates](#)
- ▶ [Module 4: Joint, Marginal, and Conditional Distributions & Functions of Random Variable](#)

Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation > Human Subjects and Special Distributions > Exponential and Uniform Distributions - Quiz

Exponential and Uniform Distributions - Quiz

🔖 Bookmark this page

Question 1

1/1 point (graded)

In the context of the exponential distribution, what is meant by memorylessness?


- ☒ a. If x describes the waiting time for some event, then the probability distribution of x at $t = 0$ is the same as the probability distribution of x at time $t = 1$ or $t = 100$ when the event has not occurred, for example. ✓
- ☐ b. The occurrence of any event does not depend on other events
- ☐ c. If x describes the waiting time for some event, then $p(x)$ follows a binomial distribution
- ☐ d. If x describes the waiting time for some event, the probability the event occurs is unrelated to the amount of time that has elapsed

Explanation


► [Module 5: Moments of a Random Variable, Applications to Auctions, & Intro to Regression](#)

▼ [Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation](#)

Human Subjects and Special Distributions

Finger Exercises due Nov 07, 2016 at 05:00 IST 

The Sample Mean, Central Limit Theorem, and Estimation

Finger Exercises due Nov 07, 2016 at 05:00 IST 

Module 6: Homework

Homework due Oct 31, 2016 at 05:00 IST 

► [Exit Survey](#)

If you think of the variable X as the wait time for some event, the probability that the event occurs at time $t = 0$ is the same as the probability that the event occurs at time $t = 10$ even if the event has not already occurred. Going back to the soccer example, if x is the time between goals, the distribution of x is the same in the 15th minute of the match as it is in the 50th minute. If the match is scoreless in the 15th minute, there is a certain probability distribution for the time until the next goal; if the match is still scoreless in the 50th minute, the probability distribution for the time until the next goal is the same.

Submit

You have used 2 of 2 attempts

✓ Correct (1/1 point)

Question 2

1 point possible (graded)

Suppose that you want to create a random variable that is exponentially-distributed in R. Which of the following methods could you use? (Select all that apply.)

- ☒ a. Use `rexp()`, which creates an exponentially-distributed random variable for you
- ☐ b. Use `runif()`, which creates a uniformly-distributed random variable, and then apply an exponential function
- ☒ c. Use `runif()` to create a uniformly-distributed random variable, then plug that into the inverse CDF formula given in class

- ☐ d. Use `runif()` to create a uniformly-distributed random variable, then plug that into the inverse PDF function



Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)

Discussion

Topic: Module 6 / Exponential and Uniform Distributions - Quiz

Show Discussion

© All Rights Reserved



© 2016 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY
OPENedX®

