



[Course](#) > [Unit 6 Linear Regression](#) > [Lecture 20: Linear Regression 2](#) > 9. Quadratic Risk and Variance

9. Quadratic Risk and Variance

Quadratic Risk





Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

The Quadratic Risk

1/1 point (graded)

We would like to analyze the *typical error in $\hat{\beta}$ compared to the true parameter, β* , which we may define as $\mathbb{E} [\|\hat{\beta} - \beta\|_2^2]$. On the other hand, we might also consider the *typical error between the predictions $\mathbb{X}\hat{\beta}$ and the observations \mathbf{Y}* , which we may define as $\mathbb{E} [\|\mathbf{Y} - \mathbb{X}\hat{\beta}\|_2^2]$.

These are respectively called the **quadratic risk of $\hat{\beta}$** and the **prediction error**. (The prediction error will be discussed in the next video.)

What happens to these errors as σ^2 increases?

☐ The error in $\hat{\beta}$ increases, but the prediction error decreases.

☐ The error in $\hat{\beta}$ decreases, but the prediction error increases.

☐ Both errors decrease.

☒ Both errors increase.



Solution:

σ^2 is the variance of each coordinate of ϵ . As the σ^2 increases, the data becomes more noisy. In particular, the task of estimating $\hat{\beta}$ ought to become harder, and it is intuitive that \mathbf{Y} becomes further from the prediction $\mathbb{X}\hat{\beta}$. To make this concrete, recall the following formulas, which hold in the homoscedastic Gaussian case:

$$\mathbb{E} [\|\hat{\beta} - \beta\|_2^2] = \sigma^2 \text{tr} ((\mathbb{X}^T \mathbb{X})^{-1})$$

$$\mathbb{E} [\|\mathbf{Y} - \mathbb{X}\hat{\beta}\|_2^2] = \sigma^2 (n - p)$$

(In our scenario, $n = 1000$ and $p = 2$.)

You have used 1 of 3 attempts

Submit

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 6 Linear Regression:Lecture 20: Linear Regression 2 / 9. Quadratic Risk and Variance

Add a Post

Show all posts ▼

by recent activity ▼

? Prediction error

2

Does it increase with more data? And decreases if there are more parameters to estimate, given the same data?

© All Rights Reserved