

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 2 Foundation of Inference](#) > [Lecture 7: Hypothesis Testing \(Continued\): Levels and P-values](#)

6. Behaviors of Type 1 and Type 2 Errors for One-Sided Tests

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

## 6. Behaviors of Type 1 and Type 2 Errors for One-Sided Tests

### How Type 1 Error Changes as Theta decreases

3/3 points (graded)

In the problems on the previous page, as well as in the examples in lecture, the level and power of the one-sided tests are determined by the type 1 and type 2 errors at the **boundary** of  $\Theta_0$  and  $\Theta_1$ . In the following problems, we will explore the qualitative reasons for this.

#### Setup:

let  $X_1, \dots, X_n \stackrel{iid}{\sim} X \sim \mathbf{P}_{\mu^*}$  where  $\mu^* \in \mathbb{R}$  is the true unknown mean of  $X$ , and the variance  $\sigma^2$  of  $X$  is fixed. The associated statistical model is  $(E, \{\mathbf{P}_{\mu}\}_{\mu \in \mathbb{R}})$  where  $E$  is the sample space of  $X$ .

We conduct a one-sided hypothesis test with the following hypotheses:

$$\begin{aligned} H_0 : \mu^* \leq \mu_0 &\iff \Theta_0 = (-\infty, \mu_0] \\ H_1 : \mu^* > \mu_0 &\iff \Theta_1 = (\mu_0, +\infty) \end{aligned}$$

Note the boundary between  $\Theta_0$  and  $\Theta_1$ . You use the statistical test:



$$\psi_n = \mathbf{1}(T_n > q_\alpha)$$


$$\text{where } T_n = \sqrt{n} \frac{\bar{X}_n - \mu_0}{\sigma}.$$

**Questions:**


Which of following regions correspond the type 1 error  $\alpha_{\psi_n}(\mu_0)$  for large  $n$ ? Note that  $\mu_0$  the boundary point of  $\Theta_0$  and  $\Theta_1$ .



(The figures on left column depicts the distribution of  $T_n$  while the ones on the right depict the distribution of  $\bar{X}_n$ . Figures not drawn to scale.)






Which orange curve below is the graph of the distribution of  $\bar{X}_n$  for  $\mu < \mu_0$ , (i.e. for  $\mu$  in the interior of  $\Theta_0$ )? The grey curve is the graph the distribution of  $\bar{X}_n$  for  $\mu = \mu_0$ .







As  $\mu$  decreases from  $\mu_0$  (i.e., moving away from the boundary of  $\Theta_0$  and  $\Theta_1$ ), does the type 1 error  $\alpha_{\psi_n}(\mu)$  increase, decrease, or not exhibit a simple trend?

☐ increase


☒ decrease

☐ does not exhibit a simple trend



### Solution:

At  $\mu = \mu_0$  and when  $n$  is large,  $T_n \sim \mathcal{N}(0, 1)$  by the CLT. Therefore, when  $n$  is large, the type 1 error  $\mathbf{P}_{\mu_0}(T_n > q_\alpha)$  is geometrically approximately the area of the "right tail" of standard normal distribution defined by the line  $T_n = q_\alpha$ .



The area of the shaded region is the type 1 error of  $\psi_n$  at  $\mu_0$ :  $\mathbf{P}_{\mu_0}(\bar{T}_n > q_\alpha)$ .


Alternatively, since

$$T_n = \sqrt{n} \frac{\bar{X}_n - \mu_0}{\sigma} > q_\alpha \iff \bar{X}_n > \mu_0 + q_\alpha \frac{\sigma}{\sqrt{n}},$$

we have

$$\mathbf{P}_{\mu_0}(T_n > q_\alpha) = \mathbf{P}_{\mu_0}\left(\bar{X}_n > \mu_0 + q_\alpha \frac{\sigma}{\sqrt{n}}\right),$$

which is the area of the "right tail" of the distribution of  $\bar{X}_n$  to the right of  $\bar{X}_n = \mu_0 + q_\alpha \frac{\sigma}{\sqrt{n}}$ . By the CLT, for  $n$  large, the distribution of  $\bar{X}_n$  is approximately Gaussian, with mean  $\mathbb{E}[X]$  and variance  $\frac{\sigma}{\sqrt{n}}$ .



The area of the shaded region is the type 1 error of  $\psi_n$  at  $\mu_0$ :  $\mathbf{P}_{\mu_0} \left( \bar{X}_n > \mu_0 + q_\alpha \frac{\sigma}{\sqrt{n}} \right)$ .

Since  $\mu = \mathbb{E}[X]$ , the CLT implies that  $\bar{X}_n$  is approximately Gaussian with mean  $\mu$  for large  $n$ . Recall the variance of  $X$  is fixed at  $\sigma$ , so the distribution of  $\bar{X}_n$  for  $\mu < \mu_0$  is a simple shift, without rescaling, to the left of the distribution of  $\bar{X}_n$  at  $\mu_0$ .


Finally, to look for a trend for the type 1 error  $\alpha_{\psi_n(\mu)}$  as  $\mu$  decreases from  $\mu_0$ , first observe that the threshold

$$\tau_{n,\alpha} = \mu_0 + q_\alpha \frac{\sigma}{\sqrt{n}}$$

of the test

$$\psi = \mathbf{1}(T_n > q_\alpha) = \mathbf{1}(\bar{X}_n > \tau_{n,\alpha})$$

does **not** depend on the parameter  $\mu$ . The only thing that changes as  $\mu$  changes is the distribution of  $\bar{X}_n$ , which shifts to the **left** as  $\mu$  decreases. Since the type 1 error  $\alpha_{\psi_n}(\mu) = \mathbf{P}_{\mu}(\bar{X}_n > \tau)$  is the area of the tail to the **right** of  $\tau$ , we see that the type 1 error continues to decrease as  $\mu$  (and the distribution of  $\bar{X}_n$ ) moves to the left.



The distribution of  $\bar{X}_n$  at  $\mu_0$ , the boundary point between  $\Theta_0$  and  $\Theta_1$ ; The distribution of  $\bar{X}_n$  at  $\mu < \mu_0$  (orange curve), a shift to the left from the distribution at  $\mu_0$

The type 1 error  $\alpha_{\psi_n}(\mu)$  in the interior of  $\Theta_0$  is smaller than the type 1 error  $\alpha_{\psi_n}(\mu_0)$  at the boundary of  $\Theta_0$  and  $\Theta_1$ .

**Remark:** The type 2 error  $\beta_{\psi_n}(\mu) = 1 - \mathbf{P}_{\mu}(\bar{X}_n > \tau)$  decreases as  $\mu$  increases from  $\mu_0$ : as  $\mu$  increases, the distribution of  $\bar{X}_n$  shifts without rescaling to the right but the threshold  $\tau$  remains constant. This implies  $\mathbf{P}_{\mu}(\bar{X}_n > \tau)$  continues to increase as  $\mu$  moves to the right from the boundary of  $\Theta_0$  and  $\Theta_1$ , and hence the Type 2 error continues to decrease.

In conclusion, for any one-sided hypothesis test where the family of distributions is parametrized by the mean of the distribution and the variance is fixed for the entire family, the type 1 and type 2 error achieve their suprema (or maxima) at the boundary between  $\Theta_0$  and  $\Theta_1$ . Therefore, the level and power can be read off at the boundary.

**Further question:** Does the reasoning above work for two-sided tests?

Submit

You have used 2 of 2 attempts



**i** Answers are displayed within the problem

## Discussion

[Hide Discussion](#)

**Topic:** Unit 2 Foundation of Inference:Lecture 7: Hypothesis Testing (Continued): Levels and P-values / 6.  
Behaviors of Type 1 and Type 2 Errors for One-Sided Tests

[Add a Post](#)

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

### **Audit Access Expires Dec 24, 2019**

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

[Learn About Verified Certificates](#)

© All Rights Reserved