



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

26:00:38



< Previous



Next >

3. Higher order partial derivatives

Bookmark this page



Calculator



Hide Notes

Lecture due Sep 13, 2021 20:30 IST Completed



Explore

Consider a function $f(x, y)$. The second partial derivative with respect to x is computed by taking the partial derivative with respect to x twice. The notation for this is

$$\frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial x^2} = (f_x)_x = f_{xx}. \quad (4.52)$$

Example 3.1 Let $f(x, y) = y^4 x^2$. Then the second partial with respect to x is given by

$$f_{xx} = \frac{\partial}{\partial x} (2y^4 x) = 2y^4. \quad (4.53)$$

We can also take second order derivatives in different variables. For example, if we first take the partial of f with respect to y and then take the partial with respect to x , we have

$$\frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) = \frac{\partial^2 f}{\partial x \partial y} = (f_y)_x = f_{yx}. \quad (4.54)$$

Example 3.2 Let $f(x, y) = y^4 x^2$. Then

$$f_{yx} = \frac{\partial}{\partial x} (4y^3 x^2) = 8y^3 x. \quad (4.55)$$

Notice the two different notations (Leibniz and subscript). Using Leibniz notation, we first take the derivative of f with respect to the variable written closest to f . So

$$\frac{\partial^2 f}{\partial x \partial y} \quad (4.56)$$

means we first take the derivative with respect to y to obtain a new function $\partial f / \partial y$. We then take the derivative of that function with respect to x . Using subscript notation, we still first take the derivative of f with respect to the variable written closest to f , but in this case, the order we write them looks reversed because the variables are on the other side of f . For example,

$$f_{xy} = \frac{\partial^2 f}{\partial y \partial x} \quad (4.57)$$

as above.

Important Fact: If a function has continuous second partial derivatives, then $f_{xy} = f_{yx}$. We explore in this course satisfy this criteria, so we will not need to worry about the order

Calculator

Hide Notes

derivatives.


Example 3.3 We saw that for $f(x,y) = y^4x^2$, we have $f_{yx} = 8y^3x$. Notice that we can also compute

$$f_{xy} = \frac{\partial}{\partial y}(2y^4x) = 8y^3x.$$

(4.58)

Higher derivatives

Start of transcript. Skip to the end.



0:00 / 0:00

2.0x

🔊

🔍

📄

🗣️

PROFESSOR: Hello, and welcome back to recitation.

The problem I like to work with you now is, simply, to compute some partial derivatives using the definitions we learned today in lecture.

So first we're going to compute the partial derivative in the x direction, of this function xy squared plus x squared y.

Video

Download video file

Transcripts

Download SubRip (.srt) file

Download Text (.txt) file

Practice 1

3/3 points (graded)
Let $u(x,t) = x^4t^2 - x^3t^5$.

Compute:

$u_{xx}(x,t) =$

12*x^2*t^2-6*x*t^5

✓ Answer: 12*x^2*t^2-6*x*t^5

$u_{tt}(x,t) =$

2*x^4-20*x^3*t^3

✓ Answer: 2*x^4-20*x^3*t^3

$u_{xt}(x,t) =$

8*x^3*t-15*x^2*t^4

✓ Answer: 8*x^3*t-15*x^2*t^4

Solution:

•

$$\begin{aligned} u_{xx}(x,t) &= \frac{\partial}{\partial x}(4x^3t^2 - 3x^2t^5) \\ &= 12x^2t^2 - 6xt^5 \end{aligned}$$

•

$$\begin{aligned}u_{tt}(x,t) &= \frac{\partial}{\partial t}(4x^3t - 5x^3t^3) \\&= 2x^4 - 20x^3t^3\end{aligned}$$

•

$$\begin{aligned}u_{xt}(x,t) &= \frac{\partial}{\partial t}(4x^3t^2 - 3x^2t^5) \\&= 8x^3t - 15x^2t^4\end{aligned}$$

Note that for the third computation, we could have computed

$$\begin{aligned}u_{tx}(x,t) &= \frac{\partial}{\partial x}(2x^4t - 5x^3t^4) \\&= 8x^3t - 15x^2t^4.\end{aligned}$$

Submit

You have used 2 of 5 attempts


 Answers are displayed within the problem

Practice 2


3/3 points (graded)
Let $g(x,y) = e^{2y} \cos(3x)$.

Compute:


$g_{xx}(x,y) =$

 **Answer:** -9*exp(2*y)*cos(3*x)

$g_{yy}(x,y) =$

 **Answer:** 4*exp(2*y)*cos(3*x)

$g_{xy}(x,y) =$

 **Answer:** -6*exp(2*y)*sin(3*x)

Solution:

•

$$\begin{aligned}g_{xx}(x,y) &= \frac{\partial}{\partial x}(-3e^{2y} \sin(3x)) \\&= -9e^{2y} \cos(3x)\end{aligned}$$

•

$$\begin{aligned}g_{yy}(x,y) &= \frac{\partial}{\partial y}(2e^{2y} \cos(3x)) \\&= 4e^{2y} \cos(3x)\end{aligned}$$

•

$$\begin{aligned}g_{xy}(x,y) &= \frac{\partial}{\partial y}(-3e^{2y} \sin(3x)) \\&= -6e^{2y} \sin(3x)\end{aligned}$$

Note that for the third computation, we could have computed

$$\begin{aligned}g_{yx}(x,y) &= \frac{\partial}{\partial x}(2e^{2y} \cos(3x)) \\&= -6e^{2y} \sin(3x).\end{aligned}$$

Submit

You have used 2 of 5 attempts

 Answers are displayed within the problem

 Calculator

 Hide Notes

 Answers are displayed within the problem

Hide Discussion

Add a Post

by recent activity ▼

- | | |
|--|---|
| <input checked="" type="checkbox"/> <u>To which textbook is speaker in video referring?</u>
<u>What textbook did they use when they taught this class live at MIT ?</u> | 7 |
| <input checked="" type="checkbox"/> <u>Geometric significance of F_{xy}.</u>
<u>F_{xx} can be thought of as a measure of how much the function bends in x direction. Is there any geometric significance like that for F_{yy}?</u> | 6 |

Next >

© All Rights Reserved



[About](#)

Affiliates

edX for Business

Open edX

Careers

[News](#)

Legal

Terms of Service & Honor Code

[Privacy Policy](#)

Accessibility Policy

Trademark Policy

 Hide Notes

[Sitemap](#)

Connect

[Blog](#)

[Contact Us](#)

[Help Center](#)

[Media Kit](#)

[Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)