



< Previous

✓

✓

Next >

8.3.3 Cooling of coffee

Bookmark this page

MO2.4

For the coffee cooling example in Section [8.2.3](#), dividing Equation ([8.16](#)) by $m_c c_c$ gives

$$\frac{dT_c}{dt} = \frac{hA}{m_c c_c} (T_{\text{out}} - T_c)$$

(8.40)

which in terms of the general IVP form gives a scalar ($M = 1$) system of equations with

$$u = T_c \quad f = \frac{hA}{m_c c_c} (T_{\text{out}} - u)$$

(8.41)

< Previous

Next >

Discussions

All posts sorted by recent activity



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)