

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Lecture 10: Consistency of MLE,
Covariance Matrices, and](#)

[Course](#) > [Unit 3 Methods of Estimation](#) > [Multivariate Statistics](#)

> 8. Covariance Matrices

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

8. Covariance Matrices

Note: Now is a good time to review the matrix exercises in [Homework 0](#).

Note on Notation: In this course, we assume all vectors to be column vectors. Therefore, while

$$\mathbf{X} = \begin{bmatrix} X^{(1)} \\ X^{(2)} \\ \vdots \\ X^{(d)} \end{bmatrix},$$

we sometimes write it as $\mathbf{X} = (X^{(1)}, \dots, X^{(d)})^T$ to be more compact in representation.

Example of Covariance II

4/4 points (graded)

Let X, Y be random variables such that

- X takes the values ± 1 each with probability 0.5
- (Conditioned on X) Y is chosen uniformly from the set $\{-3X - 1, -3X, -3X + 1\}$.

(Round all answers to 2 decimal places.)

What is $\text{Cov}(X, X)$ (equivalent to $\text{Var}(X)$)?

$\text{Cov}(X, X) =$ ✓ Answer: 1.0

What is $\text{Cov}(Y, Y)$ (equivalent to $\text{Var}(Y)$)?

$\text{Cov}(Y, Y) =$ ✓ Answer: 9.67

What is $\text{Cov}(X, Y)$?

$\text{Cov}(X, Y) =$ ✓ Answer: -3.00

What is $\text{Cov}(Y, X)$?

$\text{Cov}(Y, X) =$ ✓ Answer: -3.00

Solution:

Observe that $\mathbb{E}[X]$ and $\mathbb{E}[Y]$ are both zero, since X is uniformly distributed over $\{\pm 1\}$ and Y is uniformly distributed over the set $\{-4, -3, -2, 2, 3, 4\}$.

- $\text{Cov}(X, X)$ is the variance of X , which equals $\mathbb{E}[X^2] - \mathbb{E}[X]^2 = p + (1 - p) = 1$.
- $\text{Cov}(Y, Y)$ is the variance of Y , which equals $\mathbb{E}[Y^2] - \mathbb{E}[Y]^2 = \frac{16+9+4+4+9+16}{6} = \frac{29}{3} \approx 9.67$.

- $\text{Cov}(X, Y)$ and $\text{Cov}(Y, X)$ are always equal, by the symmetry of the definition. Observe that the joint density of (X, Y) is uniform over the pairs $(1, -4), (1, -3), (1, -2), (-1, 2), (-1, 3), (-1, 4)$. Thus, either covariance can be computed as $\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y] = \frac{-4-3-2-2-3-4}{6} = -3$.

Submit

You have used 2 of 3 attempts

i Answers are displayed within the problem

Covariance Matrix

4/4 points (graded)

Given random variables $X^{(1)}, X^{(2)}, \dots, X^{(d)}$, one can write down the **covariance matrix** Σ , where $\Sigma_{i,j} = \text{Cov}(X^{(i)}, X^{(j)})$.

Let $X^{(1)}, X^{(2)}$ be random variables such that

- $X^{(1)}$ takes the values ± 1 each with probability 0.5
- (Conditioned on $X^{(1)}$) $X^{(2)}$ is chosen uniformly from the set $\{-3X^{(1)} - 1, -3X^{(1)}, -3X^{(1)} + 1\}$.

What is the covariance matrix Σ ?

$$\Sigma_{1,1} = \boxed{1} \quad \checkmark \text{ Answer: } 1.0 \quad \Sigma_{1,2} = \boxed{-3} \quad \checkmark \text{ Answer: } -3.00$$

$$\Sigma_{2,1} = \boxed{-3} \quad \checkmark \text{ Answer: } -3.00 \quad \Sigma_{2,2} = \boxed{29/3} \quad \checkmark \text{ Answer: } 9.67$$

Solution:

Using the answer to the previous question, the 2×2 covariance matrix Σ evaluates to

$$\Sigma = \begin{pmatrix} \Sigma_{1,1} & \Sigma_{1,2} \\ \Sigma_{2,1} & \Sigma_{2,2} \end{pmatrix} = \begin{pmatrix} 1 & -3 \\ -3 & \frac{29}{3} \end{pmatrix}$$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Covariance Matrix: Definitions



I have d rows and d columns.

5:11 / 5:11

1.50x

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Here is a compact formula for the covariance matrix using vector notation.

Let $\mathbf{X} = \begin{pmatrix} X^{(1)} \\ \vdots \\ X^{(d)} \end{pmatrix}$ be a random vector of size $d \times 1$.

Let $\mu \triangleq \mathbb{E}[\mathbf{X}]$ denote the **entry-wise** mean, i.e.

$$\mathbb{E}[\mathbf{X}] = \begin{pmatrix} \mathbb{E}[X^{(1)}] \\ \vdots \\ \mathbb{E}[X^{(d)}] \end{pmatrix}.$$

Consider the vector outer product (refer to [Homework 0](#)) $(\mathbf{X} - \mu)(\mathbf{X} - \mu)^T$, which is a random $d \times d$ matrix. Then the **covariance matrix** Σ can be written as

$$\Sigma = \mathbb{E}[(\mathbf{X} - \mu)(\mathbf{X} - \mu)^T].$$

Note: The following exercises will be discussed as properties of covariance in the video that follows, but we encourage you attempt these exercises before watching the video.

Covariance Matrix: Properties I

1/1 point (graded)

Let \mathbf{X} be a random vector and let $\mathbf{Y} = \mathbf{X} + B$, where B is a constant vector. Let $\mu_{\mathbf{X}}$ be the mean vector of \mathbf{X} and let $\Sigma_{\mathbf{X}}$ be the covariance matrix of \mathbf{X} . Select from the following all statements that are correct.

☐ The covariance matrix of \mathbf{Y} could potentially be equal to $\Sigma_{\mathbf{X}}$ only under some conditions imposed on B

☒ The covariance matrix of \mathbf{Y} is the same as $\Sigma_{\mathbf{X}}$ for all vectors B

☒ The covariance matrix of \mathbf{Y} has the same size as the matrix $\Sigma_{\mathbf{X}}$

☐ The covariance matrix of \mathbf{Y} is the same as $\Sigma_{\mathbf{X}}$ if and only if vector B is equal to 0



Solution:

Choices 2 and 3 are correct. Let the covariance matrix of \mathbf{Y} be denoted $\Sigma_{\mathbf{Y}}$. Note that $\mathbb{E}[\mathbf{X} + B] = \mu_{\mathbf{X}} + B$ for any vector B .

$$\Sigma_{\mathbf{Y}} = \mathbb{E}[(\mathbf{X} + B - \mu_{\mathbf{X}} - B)(\mathbf{X} + B - \mu_{\mathbf{X}} - B)^T] = \Sigma_{\mathbf{X}}$$

Since choice 2 is correct, choices 1 and 4 that impose certain conditions on B are technically incorrect as we do not require that B satisfy some conditions for $\Sigma_{\mathbf{Y}}$ to be the same as $\Sigma_{\mathbf{X}}$.

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Covariance Matrix: Properties II

1/1 point (graded)

Let \mathbf{X} be a random vector of size $d \times 1$ and let $\mathbf{Y} = A\mathbf{X} + B$, where A is a constant matrix of size $n \times d$ and B is a constant vector of size $n \times 1$. Let $\mu_{\mathbf{X}}$ be the mean vector of \mathbf{X} and let $\Sigma_{\mathbf{X}}$ be the covariance matrix of \mathbf{X} . Let $\mu_{\mathbf{Y}}$ be the mean vector of \mathbf{Y} and let $\Sigma_{\mathbf{Y}}$ be the covariance matrix of \mathbf{Y} .

Select from the following all statements that are correct.

☒ $\Sigma_{\mathbf{Y}}$ is the same as covariance matrix of $A\mathbf{X}$

☒ $\Sigma_{\mathbf{Y}}$ is of size $n \times n$

☐ $\Sigma_{\mathbf{Y}} = A^2 \Sigma_{\mathbf{X}}$

☒ $\Sigma_{\mathbf{Y}} = A \Sigma_{\mathbf{X}} A^T$

☐ $\Sigma_{\mathbf{Y}} = A^T \Sigma_{\mathbf{X}} A$



Solution:

As \mathbf{Y} is an $n \times 1$ random vector, $\Sigma_{\mathbf{Y}}$ is of size $n \times n$.

From the previous problem we know that $\Sigma_{\mathbf{Y}}$ is the same as the covariance matrix of $A\mathbf{X}$. Therefore, it suffices to find this matrix, which we denote $\Sigma_{A\mathbf{X}}$.

$$\begin{aligned}
 \Sigma_{A\mathbf{X}} &= \mathbb{E} \left[(A\mathbf{X} - A\mu_X) (A\mathbf{X} - A\mu_X)^T \right] \\
 &= \mathbb{E} \left[A(\mathbf{X} - \mu_X) (\mathbf{X}^T A^T - \mu_X^T A^T) \right] \\
 &= \mathbb{E} \left[A(\mathbf{X} - \mu_X) (\mathbf{X} - \mu_X)^T A^T \right] \\
 &= A \mathbb{E} \left[(\mathbf{X} - \mu_X) (\mathbf{X} - \mu_X)^T \right] A^T \\
 &= A \Sigma_{\mathbf{X}} A^T.
 \end{aligned}$$

Therefore, choices 1, 2, and 4 are correct.

Choices 3 and 5 are not correct in general (even if A is a square matrix) because matrix multiplication is not commutative.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Covariance Matrix: Affine Transformation

The video content shows a chalkboard with the following derivations:

$$\begin{aligned} E(XY) &= E(X)E(Y) + \text{Cov}(X, Y) \\ &= E(X)E(Y) + \text{Cov}(X, Y) \\ &= E(X)E(Y) + \text{Cov}(X, Y) \end{aligned}$$

Below the chalkboard, a lecturer is visible. The video player interface shows a play button in the center, a progress bar at the bottom left indicating 6:18 / 6:18, and control icons at the bottom right for volume, full screen, and subtitles.

Video

Transcripts

[Download video file](#)[Download SubRip \(.srt\) file](#)[Download Text \(.txt\) file](#)

Effect of Linear Transformations of Covariance Matrix

4/4 points (graded)

Let $\mathbf{X} = \begin{pmatrix} X^{(1)} \\ X^{(2)} \end{pmatrix}$ be a random vector with covariance Matrix $\Sigma_{\mathbf{X}} = \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}$.

Let $\mathbf{Y} = M\mathbf{X}$, where $M = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$.

Observe that $Y^{(1)} = X^{(1)} - X^{(2)}$ and $Y^{(2)} = X^{(1)} + X^{(2)}$. What is the new covariance matrix $\Sigma_{\mathbf{Y}}$?

$(\Sigma_{\mathbf{Y}})_{1,1} =$ ✓ Answer: 1 $(\Sigma_{\mathbf{Y}})_{1,2} =$ ✓ Answer: 0.0

$(\Sigma_{\mathbf{Y}})_{2,1} =$ ✓ Answer: 0.0 $(\Sigma_{\mathbf{Y}})_{2,2} =$ ✓ Answer: 3

Solution:

Recall from an earlier problem that for any pair of random variables A, B with the same variance $\text{Var}(A) = \text{Var}(B) = \sigma^2$, $\text{Cov}(A - B, A + B) = \text{Var}(A) - \text{Var}(B) = 0$.

Therefore, given the matrix M , $\Sigma_{\mathbf{Y}}$ must be a diagonal matrix.

We have

$$\text{Cov}(Y^{(1)}, Y^{(1)}) = \text{Cov}(X^{(1)} - X^{(2)}, X^{(1)} - X^{(2)}) = \text{Cov}(X^{(1)}, X^{(1)}) - 2\text{Cov}(X^{(1)}, X^{(2)}) + \text{Cov}(X^{(2)}, X^{(2)}) = 1 - 1 + 1 = 1.$$

Similarly,

$$\text{Cov}(Y^{(2)}, Y^{(2)}) = \text{Cov}(X^{(1)} + X^{(2)}, X^{(1)} + X^{(2)}) = \text{Cov}(X^{(1)}, X^{(1)}) + 2\text{Cov}(X^{(1)}, X^{(2)}) + \text{Cov}(X^{(2)}, X^{(2)}) = 1 + 1 + 1 = 3.$$

You have used 2 of 3 attempts

i Answers are displayed within the problem

Discussion

[Hide Discussion](#)

Topic: Unit 3 Methods of Estimation: Lecture 10: Consistency of MLE, Covariance Matrices, and Multivariate Statistics / 8. Covariance Matrices

[Add a Post](#)

Show all posts ▼

by recent activity ▼

- ? Variance tells us variation of a r.v., Covariance tells us variation b/w 2 r.v. is there [Something] that tells us about variation of m r.v 13
- 💬 The last problem seems simpler to just solve... 3
- 💬 One of the Cov(A, A) questions seems to be done in terms of X and then cancelling out X
But keep getting this marked as incorrect. Is there a way to compute this, or should I also iterate through the values of X. This for Question... **Example of Covariance II** 2
- ✓ Example of Covariance II
In the first exercise I can't get cov(x,y) right although I did get cov(x,x) and cov(y,y) correct so not sure what may be wrong? Once I displayed all possible x,y pairs in an excel sh... 5

[Learn About Verified Certificates](#)

© All Rights Reserved