



Bookmarks

- ▶ Unit 0: Overview
- ▶ Entrance Survey
- ▶ Unit 1: Probability models and axioms
- ▶ Unit 2: Conditioning and independence
- ▶ Unit 3: Counting
- ▶ Unit 4: Discrete random variables
- ▶ Exam 1
- ▶ Unit 5: Continuous random variables

Exam 2 > Exam 2 > Exam 2 vertical3

Bookmark

Problem 4: Manhole covers

(2/3 points)

Manhole explosions (usually caused by gas leaks and sparks) are on the rise in your city. On any given day, the manhole cover near your house explodes with some unknown probability, which is the same across all days. We model this unknown probability of explosion as a random variable Q , which is uniformly distributed between **0** and **0.1**. Let X_i be a Bernoulli random variable that indicates whether the manhole cover near your house explodes on day i (where today is day **1**).

Give numerical answers for parts (1) and (2).

1.

 $\mathbf{E}[X_i] =$ Answer: 0.05

2.

 $\mathbf{var}(X_i) =$ Answer: 0.0475

► Unit 6: Further topics on random variables

► Unit 7: Bayesian inference

▼ Exam 2

Exam 2

Exam 2 due Apr 20, 2016 at 23:59 UTC



► Unit 8: Limit theorems and classical statistics

3. Let \mathbf{A} be the event that the manhole cover did not explode yesterday (i.e., $X_0 = 0$). Find the conditional PDF of Q given \mathbf{A} . Express your answer in terms of q using standard notation .

$$10/(1-5*q^2)$$

✗

For $0 \leq q \leq 0.1$, $f_{Q|\mathbf{A}}(q) =$ Answer: $200*(1-q)/19$

Answer:

1. Using the law of iterated expectations, we have

$$\mathbf{E}[X_i] = \mathbf{E}[\mathbf{E}[X_i | Q]] = \mathbf{E}[Q] = 0.05.$$

2. Using the law of total variance, we have

$$\begin{aligned} \text{var}(X_i) &= \mathbf{E}[\text{var}(X_i | Q)] + \text{var}(\mathbf{E}[X_i | Q]) \\ &= \mathbf{E}[Q(1 - Q)] + \text{var}(Q) \\ &= \mathbf{E}[Q] - \mathbf{E}[Q^2] + \text{var}(Q) \\ &= \mathbf{E}[Q] - (\text{var}(Q) + (\mathbf{E}[Q])^2) + \text{var}(Q) \\ &= \mathbf{E}[Q] - (\mathbf{E}[Q])^2 \\ &= 0.05 - 0.05^2 \\ &= 0.0475. \end{aligned}$$

3. Using Bayes' rule, we have for $0 \leq q \leq 0.1$,

$$\begin{aligned} f_{Q|A}(q) &= \frac{f_Q(q)\mathbf{P}(A \mid Q = q)}{\mathbf{P}(A)} \\ &= \frac{f_Q(q)\mathbf{P}(A \mid Q = q)}{\int_0^{0.1} f_Q(q)\mathbf{P}(A \mid Q = q) dq} \\ &= \frac{10(1 - q)}{\int_0^{0.1} 10(1 - q) dq} \\ &= \frac{10(1 - q)}{1 - 0.05} \\ &= \frac{200(1 - q)}{19}. \end{aligned}$$

You have used 4 of 4 submissions

© All Rights Reserved



© edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

