



[Course](#) > [Unit 4 Hypothesis testing](#) > [T-Test](#)

8. Sample Variance and Sample  
Mean of IID Gaussians: Cochran's  
> Theorem

**Audit Access Expires Dec 24, 2019**

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 8. Sample Variance and Sample Mean of IID Gaussians: Cochran's Theorem

### Cochran's Theorem: Independence of Gaussian Sample Variance and Sample Mean

## Important example: the sample variance

- ▶ Recall that the sample variance is given by

$$S_n = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - (\bar{X}_n)^2$$

- ▶ Cochran's theorem states that for  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ , if

- ▶  $\bar{X}_n \perp\!\!\!\perp S_n$ ; for all  $n$ .

- ▶  $\frac{nS_n}{\sigma^2} \sim \chi_{n-1}^2$ .

$$E[S_n] = \frac{n-1}{n} \sigma^2$$

- ▶ We often prefer the unbiased estimator of  $\sigma^2$ :

$$\tilde{S}_n = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = \frac{n}{n-1} S_n$$

$$E[\tilde{S}_n] = \frac{n}{n-1} E\left[\frac{\sigma^2}{n} \chi_{n-1}^2\right] = \frac{n\sigma^2}{n-1} \cdot \frac{n-1}{n} = \sigma^2$$

▶ 7:29 / 7:29

▶ 1.50x



### Video

[Download video file](#)

### Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

## A Special Case of Cochran's Theorem I

3/3 points (graded)

**Cochran's theorem** states that if  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ , then the sample variance

$$S_n := \frac{1}{n} \left( \sum_{i=1}^n X_i^2 \right) - (\bar{X}_n)^2$$

satisfies:

- $\bar{X}_n$  is independent of  $S_n$ , and
- $\frac{nS_n}{\sigma^2} \sim \chi_{n-1}^2$ .

In this problem, you will verify that Cochran's theorem holds when  $n = 2$ . Let  $X_1, X_2 \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ .

The expression  $S_2$  can be written in the form  $A^2$  where  $A$  is a polynomial in  $X_1$  and  $X_2$ .

What is  $A^2$ ?

Type **X\_1** for  $X_1$  and **X\_2** for  $X_2$ .

$$A^2 = \boxed{(X_1 - X_2)^2 / 4} \quad \checkmark \text{ Answer: } (X_1 - X_2)^2 / 4$$

The expression  $A$  from the previous question is a random variable, and moreover is distributed as  $\mathcal{N}(\mu^*, (\sigma^*)^2)$  for some  $\mu^*$  and  $\sigma^*$  that can be expressed in terms of the original parameters  $\mu$  and  $\sigma$ . (Note:  $A$  can have two forms, but both would have the same distribution by symmetry).

What is  $\mu^*$  expressed in terms of  $\mu$  and  $\sigma$ ?

$$\mu^* = \boxed{0} \quad \checkmark \text{ Answer: } 0.0$$

What is  $(\sigma^*)^2$  expressed in terms of  $\mu$  and  $\sigma$ ?

$$(\sigma^*)^2 = \boxed{\text{sigma}^2/2} \quad \checkmark \text{ Answer: sigma}^2/2$$

$$\frac{\sigma^2}{2}$$

STANDARD NOTATION

**Solution:**

Observe that

$$S_n = \frac{X_1^2 + X_2^2}{2} - \left( \frac{X_1 + X_2}{2} \right)^2 = \frac{X_1^2}{4} + \frac{X_2^2}{4} - \frac{1}{2}X_1X_2 = \left( \frac{X_1 - X_2}{2} \right)^2.$$

Hence, we can take  $A = \pm \frac{X_1 - X_2}{2}$  (either choice has the same distribution, by symmetry). Next,

$$\mathbb{E}[A] = \frac{1}{2}\mathbb{E}[X_1 - X_2] = \frac{1}{2}(\mu - \mu) = 0,$$

and

$$\text{Var}(A) = \text{Var}\left(\frac{X_1 - X_2}{2}\right) = \frac{1}{4}(\text{Var}(X_1) + \text{Var}(X_2)) = \frac{\sigma^2}{2}.$$

Submit

You have used 2 of 3 attempts

**i** Answers are displayed within the problem

A Special Case of Cochran's Theorem II

4/4 points (graded)

As above, let  $X_1, X_2 \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ .

Recall the random variable  $A$  that you found in the previous problem in terms of  $X_1$  and  $X_2$ .

Let  $\bar{X}_2 = \frac{X_1 + X_2}{2}$ , i.e.  $\bar{X}_n$  when  $n = 2$ .

What is  $\mathbb{E}[A\bar{X}_2]$ ?

0

✓ Answer: 0.0

Using the answer above, which of the following are true? (Choose all that apply.)

☒  $A$  and  $\bar{X}_2$  are independent.

☐  $A$  and  $\bar{X}_2$  are not independent.

☐  $A, \bar{X}_2 \sim \mathcal{N}(0, 2\sigma^2)$ .

☒  $A \sim \mathcal{N}(0, \sigma^2/2)$  and  $\bar{X}_2 \sim \mathcal{N}(\mu, \sigma^2/2)$ .



For some expression  $B$  in terms of  $\sigma^2$ , the random variable  $BS_2 \sim \chi^2$ . What is  $B$ ?

$B =$

2/sigma^2

✓ Answer: 2/sigma^2

$\frac{2}{\sigma^2}$

STANDARD NOTATION

How many degrees of freedom does the  $\chi^2$  random variable  $BS_2$  have?

✓ Answer: 1

**Solution:**

Recall that  $A = \frac{X_1 - X_2}{2}$  and  $\bar{X}_2 = \frac{X_1 + X_2}{2}$ . Hence,

$$\mathbb{E}[A\bar{X}_2] = \frac{1}{4}\mathbb{E}[(X_1 - X_2)(X_1 + X_2)] = \frac{1}{4}(\sigma^2 - \sigma^2) = 0.$$

As jointly Gaussian variables (why is it that  $A$  and  $\bar{X}_2$  are jointly Gaussian?) that are uncorrelated are also independent,  $A$  and  $\bar{X}_2$  are independent. By the previous problem, we know  $A \sim \mathcal{N}(0, \sigma^2/2)$ . A quick calculation shows that  $\bar{X}_2 \sim \mathcal{N}(\mu, \sigma^2/2)$ . Hence, the first and last choices are correct in the multiple choice question.

Observe that

$$\frac{2}{\sigma^2}S_2 = \frac{2}{\sigma^2}\left(\frac{X_1 - X_2}{2}\right)^2 = \left(\frac{X_1 - X_2}{\sqrt{2}\sigma}\right)^2,$$

and  $\frac{X_1 - X_2}{\sqrt{2}\sigma} \sim \mathcal{N}(0, 1)$ . By definition,  $\frac{2}{\sigma^2}S_2 \sim \chi_1^2$ .

**Remark:** The last question shows that  $\frac{2}{\sigma^2}S_2 \sim \chi_1^2$ , which verifies the second claim in Cochran's theorem for this special case. To show the first part of Cochran's theorem, that  $S_2$  and  $\bar{X}_2$  are independent, recall that we showed  $A = \sqrt{S_2}$  is independent of  $\bar{X}_2$ . By a standard fact of probability, this also implies that  $A^2 = S_2$  is independent of  $\bar{X}_2$ .

Submit

You have used 1 of 3 attempts

## Concept Check: Cochran's Theorem and Unbiased Sample Variance

1/1 point (graded)

Let  $X_1, \dots, X_n$  be i.i.d. and distributed according to  $\mathcal{N}(0, \sigma^2)$ . Let  $\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$ . What is the distribution of  $\frac{(n-1)\tilde{S}_n}{\sigma^2}$ , where  $\tilde{S}_n$  is the unbiased sample variance of  $X_1, \dots, X_n$ :

$$\tilde{S}_n = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2$$

Type **Cn** for chi-squared distribution with  $n$  degrees of freedom, **Cn1** for chi-squared distribution with  $n - 1$  degrees of freedom.

Cn1

✓ Answer: Cn1 + 0\*Cn

Cn1

STANDARD NOTATION

### Solution:

By Cochran's theorem,

$$\begin{aligned} \frac{nS_n}{\sigma^2} &\sim \chi_{n-1}^2 \\ \Leftrightarrow \frac{(n-1)\tilde{S}_n}{\sigma^2} &\sim \chi_{n-1}^2 \end{aligned}$$

**Remark:** We will use the random variable  $\frac{\tilde{S}_n}{\sigma^2}$  in the upcoming videos in what is called the Student's T Test. The point of this problem was to show that  $\frac{(n-1)\tilde{S}_n}{\sigma^2}$  is a  $\chi_{n-1}^2$  random variable, thereby showing that the distribution of  $\frac{\tilde{S}_n}{\sigma^2}$  is the distribution of a  $\chi_{n-1}^2$  random variable scaled by  $\frac{1}{n-1}$ .

Submit

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Discussion


Hide Discussion

**Topic:** Unit 4 Hypothesis testing:Lecture 13: Chi Squared Distribution, T-Test / 8. Sample Variance and Sample Mean of IID Gaussians: Cochran's Theorem

Add a Post

Show all posts ▼

by recent activity ▼

 [Deep Dive/Recitation on Chi-Squared, Students t Distributions, and Independence of the Sample Mean and Variance for iid Normal Variables](#)

1

 [Pinned](#)  [Community TA](#)

Learn About Verified Certificates

© All Rights Reserved