

<u>Course</u> > <u>Midterm Exam 1</u> > <u>Midterm Exam 1</u> > 2.

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

2. Setup:

The lifetime (in months) of a battery is modeled by a random variable X that has pdf

$$f_{ heta}\left(x
ight)=K heta^{x}\mathbf{1}\left(x>0
ight) \qquad ext{where }K=\ln\left(1/ heta
ight)$$

for an unknown parameter $\theta \in (0,1)$. (Here $\mathbf{1}$ (x>0) is the indicator variable that takes value 1 when its argument is true, i.e. when x>0.)

Assume that we have n independent observations X_1,\ldots,X_n of the lifetime of n batteries. We want to use these observations to estimate $\theta\in(0,1)$.

Note (October 24): Assume that the observations are of the same type of batteries.

Input instructions: For all problems below, when inputting the natural log function, use \ln , e.g. enter $\ln(\text{theta})$ for $\ln(\theta)$.

Expectation and Variance

2/2 points (graded)

Compute the expected value $\mathbb{E}\left[X_{i}
ight]$ and the variance $\mathsf{Var}\left[X_{i}
ight]$ of X_{i} .

(Enter your answer in terms of θ **only**.)

Hint: Note that the given pdf is equivalent to the pdf of a common distribution except reparametrized with a different parameter.

$$\mathbb{E}\left[X_i\right] = \boxed{-1/\ln(\text{theta})}$$

$$-\frac{1}{\ln(\theta)}$$

$$\text{Var}\left[X_i\right] = \boxed{1/(\ln(\text{theta}))^2}$$

$$\frac{1}{(\ln(\theta))^2}$$

$$\text{STANDARD NOTATION}$$

Solution:

f is the pdf of an exponential reparametrized in the paramter heta instead of the usual λ where $heta=e^{-\lambda}$, or equivalently $\lambda=\ln{(1/ heta)}$. Hence, $X_i\sim \mathsf{Exp}\left(\ln{(1/ heta)}\right)$ and

$$\mathbb{E}\left[X_i
ight] \;=\; rac{1}{\lambda} \;=\; \left(\ln\left(1/ heta
ight)
ight)^{-1}$$
 $\mathsf{Var}\left[X_i
ight] \;=\; rac{1}{\lambda^2} \;=\; \left(\ln\left(1/ heta
ight)
ight)^{-2}.$

Submit

You have used 1 of 3 attempts

1 Answers are displayed within the problem

Maximum Likelihood Estimator

3.0/3.0 points (graded)

Compute the maximum likelihood estimator $\hat{\theta}$ of θ .

(Enter $extbf{barX_n}$ for the sample average $ar{X}_n = rac{1}{n} \sum_{i=1}^n X_i$.)

$$\hat{\theta} = \left| \begin{array}{c} e^{-1/barX_n} \end{array} \right|$$

✓ Answer: e^(-(1/barX_n))

STANDARD NOTATION

Submit

You have used 1 of 3 attempts

• Answers are displayed within the problem

Asymptotic Distribution

1/1 point (graded)

What kind of distribution does $\sqrt{n}\,(\hat{ heta}- heta)$ converge in distribution to?



nonparametric





unknown

~

Submit

You have used 1 of 3 attempts

• Answers are displayed within the problem

Asymptotic Variance

3.0/3.0 points (graded)

Find the asymptotic variance $V\left(\theta\right)$ of $\hat{\theta}$.

STANDARD NOTATION

Solution:

Use either the Fisher information or the Delta Method. $I\left(heta
ight)=rac{1}{ heta^{2}(\ln heta)^{2}}$

Submit

You have used 1 of 3 attempts

• Answers are displayed within the problem

Confidence Interval

2.0/2.0 points (graded)

Find the one-sided confidence interval of the form $\mathcal{I}_{ ext{plug-in}}=(0,B]$ at asymptotic (confidence) level 95% for θ given by the plug-in method. Specify B below in terms of $n,\hat{\theta}$ and $V(\hat{\theta})$.

(Enter **hattheta** for $\hat{\theta}$, **V** for $V(\hat{\theta})$.

If applicable, for any numerical α , enter **q(alpha)** for the $1-\alpha$ quantile q_{α} of the standard normal distribution, i.e. $\mathbf{P}(Z \leq q_{\alpha}) = 1-\alpha)$. For, example enter **q(0.01)** for $q_{0.01}$;

Note (Oct 19): Earlier verion contained an error in the input instructions, and stated "If applicable, for any numerical α , enter **q(alpha)** for the $1-\alpha$ quantile q_{α} of the standard normal distribution, i.e. $\mathbf{P}\left(Z\leq q_{\alpha}\right)=\alpha$)." You may also use numerical values for q_{α} as long as they are accurate to at least 4 decimal places.)

$$B = \left| \text{hattheta+q(0.05)*sqrt(V/n)} \right|$$

✓ Answer: hattheta+q(0.05)*sqrt(V/n)

STANDARD NOTATION

Submit

You have used 1 of 3 attempts

• Answers are displayed within the problem

Confidence Interval Continued

1/1 point (graded)

We observe $\hat{ heta}=0.62$ for n=100 .

(Give an answer accurate to at least 3 decimal places.)

$$B = 0.6687505331016123$$

✓ Answer: 0.6687

STANDARD NOTATION

Submit

You have used 1 of 3 attempts


1 Answers are displayed within the problem

Error and Bug Reports/Technical Issues

Topic: Midterm Exam 1:Midterm Exam 1 / 2.

Hide Discussion

Add a Post



Confidence Interval		3	
? [Staff] removed by staff		2	•
	Learn About Verified Certificates		
	Zearry to dat Vermed Cerementes	© All Ri	ights Reserved