EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the <u>Privacy Policy</u>.

×



Lecture 10: Consistency of MLE, Covariance Matrices, and

<u>Course</u> > <u>Unit 3 Methods of Estimation</u> > <u>Multivariate Statistics</u>

> 9. Multivariate Gaussian Distribution

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.


Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

9. Multivariate Gaussian Distribution

Note: Now is a good time to review Gaussian random variables from <u>Lecture 2</u>.

Video Note: In the slide of the video below, there is a typo in the formula of the pdf of the multivariate Gaussian distribution: the exponent d in overall scaling factor should apply only to 2π , rather than $2\pi \det \Sigma$. The correct version is in the note below the video. (The unannotated slides in the resource section have also been corrected).

Multivariate Gaussian Distribution: Definition



Video

<u>Download video file</u>

Transcripts

<u>Download SubRip (.srt) file</u>

<u>Download Text (.txt) file</u>

Multivariate Gaussian Random Variable

A random vector $\mathbf{X} = \left(X^{(1)}, \dots, X^{(d)}\right)^T$ is a **Gaussian vector**, or **multivariate Gaussian or normal variable**, if any linear combination of its components is a (univariate) Gaussian variable or a constant (a "Gaussian" variable with zero variance), i.e., if $\alpha^T \mathbf{X}$ is (univariate) Gaussian or constant for any constant non-zero vector $\alpha \in \mathbb{R}^d$.

The distribution of \mathbf{X} , the d-dimensional Gaussian or normal distribution , is completely specified by the vector mean $\mu = \mathbb{E}\left[\mathbf{X}\right] = \left(\mathbb{E}\left[X^{(1)}\right], \dots, \mathbb{E}\left[X^{(d)}\right]\right)^T$ and the $d \times d$ covariance matrix Σ . If Σ is invertible, then the pdf of \mathbf{X} is

$$f_{\mathbf{X}}\left(\mathbf{x}
ight) = rac{1}{\sqrt{\left(2\pi
ight)^{d}\mathrm{det}\left(\Sigma
ight)}}e^{-rac{1}{2}\left(\mathbf{x}-\mu
ight)^{T}\Sigma^{-1}\left(\mathbf{x}-\mu
ight)}, \;\;\; \mathbf{x} \in \mathbb{R}^{d}$$

where $\det\left(\Sigma\right)$ is the determinant of the Σ , which is positive when Σ is invertible.

If $\mu=\mathbf{0}$ and Σ is the identity matrix, then \mathbf{X} is called a **standard normal random vector** .

Note that when the covariant matrix Σ is diagonal, the pdf factors into pdfs of univariate Gaussians, and hence the components are independent.

Linear Transformation of a Multivariate Gaussian Random Vector

1/1 point (graded)

Consider the 2-dimensional Gaussian
$$\mathbf{X}=egin{pmatrix} X^{(1)} \ X^{(2)} \end{pmatrix}$$
 with covariance matrix $\Sigma_X=egin{pmatrix} 1 & 2 \ 2 & 5 \end{pmatrix}$ and mean $\mu_{\mathbf{X}}=egin{pmatrix} 0 \ 0 \end{pmatrix}$.

Consider the vector $lpha=inom{1}{-1}$, so that $Y=lpha^T\mathbf{X}$ is a 1-dimensional Gaussian.

What is the variance Var(Y) of Y?

$$Var(Y) =$$
 2 Answer: 2

Solution:

One way to answer this is to notice that $Y=X^{(1)}-X^{(2)}$, so

$$\mathsf{Var}\left(Y\right) = \mathsf{Cov}\left(Y,Y\right) = \mathsf{Var}\left(X^{(1)}\right) + \mathsf{Var}\left(X^{(2)}\right) - 2\mathsf{Cov}\left(X^{(1)},X^{(2)}\right) = 1 + 5 - 4 = 2.$$

Another way is to define the matrix $M \triangleq \alpha^T = (1 \quad -1)$, and apply the formula $\Sigma_Y = M \Sigma_{\mathbf{X}} M^T = 2$.

Submit

You have used 2 of 3 attempts

Answers are displayed within the problem

Singular Covariance Matrices

1/1 point (graded)

Consider again a 2-dimensional Gaussian
$$\mathbf{X} = \begin{pmatrix} X^{(1)} \\ X^{(2)} \end{pmatrix}$$
. But instead, Σ_X is $\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$ and $\alpha = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$, what is the variance $\mathsf{Var}(Y)$ of $Y = \alpha^T \mathbf{X}$?

This result tells us that the Gaussian $(X^{(1)}, X^{(2)})^T$ is actually a one-dimensional Gaussian, orthogonal to the direction of $\begin{pmatrix} 2 \\ -1 \end{pmatrix}$.

Solution:

Define a matrix $M=lpha^T$. We have $\Sigma_Y=M\Sigma_XM^T=0$, since M^T is a column vector in the nullspace of Σ_X .

Such a Gaussian (with a singular covariance matrix) is sometimes referred to as a **degenerate** Gaussian.

Submit

You have used 1 of 3 attempts

• Answers are displayed within the problem

(Optional) Diagonalization of the Covariance Matrix

Show

(Optional) Gaussian Random Vectors I

0 points possible (ungraded)

Recall from an earlier part of this lecture that the covariance between two random variables being 0 does not necessarily imply that the random variables are independent. However, this is true if the random variables are multivariate Gaussian.

Let X be a Gaussian random vector with mean μ and covariance Σ . Assume that Σ is positive definite. Determine if the following statement is true or false.

"There exists a vector B and a matrix A such that $A(\mathbf{X} + B)$ is a Gaussian random vector whose components are independent and each of mean 0".







Hint: Refer to the note above on diagonalization of the covariance matrix.

Solution:

True. First, in order to remove the effect of μ we can set $B=-\mu$ to make the individual Gaussian random variables be of zero mean. Let $\widehat{\mathbf{X}}=\mathbf{X}-\mu$. From an earlier problem we know that the covariance matrix of $\widehat{\mathbf{X}}$ is the same as Σ .

From the above note on covariance matrices we can see that there exists an orthogonal matrix U such that $D=U\Sigma U^T$.

Consider the following transformation: $\mathbf{Y} = U\widehat{\mathbf{X}}$.

The covariance matrix of \mathbf{Y} is (from an earlier problem)

$$U\Sigma U^T$$
,

which is precisely equal to the diagonal matrix D. Therefore, \mathbf{Y} has component Gaussian random variables that are uncorrelated and hence independent.

Submit

You have used 1 of 1 attempt

• Answers are displayed within the problem

Discussion

Topic: Unit 3 Methods of Estimation:Lecture 10: Consistency of MLE, Covariance Matrices, and Multivariate Statistics / 9. Multivariate Gaussian Distribution

Add a Post

Hide Discussion



Learn About Verified Certificates

© All Rights Reserved