

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



Homework 4: TV distance, KL-

Course > Unit 3 Methods of Estimation > Divergence, and Introduction to MLE > 2. Compute Total Variation Distance

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

2. Compute Total Variation Distance

(a)


3/3 points (graded)

Compute the total variation distance between

$$\mathbf{P} = X \quad \text{and} \quad \mathbf{Q} = X + c, \quad \text{where } X \sim \text{Ber}(p), p \in (0, 1), \text{ and } c \in \mathbb{R}.$$

(If applicable, enter **abs(x)** for $|x|$. Simplify your answer to have the minimum number of absolute signs possible.)

For $c \notin \{-1, 0, 1\}$:

$\text{TV}(\mathbf{P}, \mathbf{Q}) =$ 

For $c = 0$:

$$\text{TV}(\mathbf{P}, \mathbf{Q}) = \boxed{0} \quad \checkmark$$

0

For $c = 1$ or $c = -1$:

$$\text{TV}(\mathbf{P}, \mathbf{Q}) = \boxed{(1+\text{abs}(1-2\cdot p))/2} \quad \checkmark$$

$\frac{1+\text{abs}(1-2\cdot p)}{2}$

STANDARD NOTATION

Submit

You have used 1 of 2 attempts

✓ Correct (3/3 points)

(b)

2/2 points (graded)

Compute the total variation distance between

$$\mathbf{P} = \text{Ber}(p) \quad \text{and} \quad \mathbf{Q} = \text{Ber}(q), \quad \text{where } p, q \in [0, 1].$$

(If applicable, enter **abs(x)** for $|x|$.)

$$\text{TV}(\mathbf{P}, \mathbf{Q}) = \boxed{\text{abs}(p-q)} \quad \checkmark$$

$\text{abs}(p - q)$

Let X_1, \dots, X_n be n i.i.d. Bernoulli random variables with some parameter $p \in [0, 1]$, and \bar{X}_n be their empirical average. Consider the total variation distance $\text{TV}(\text{Ber}(\bar{X}_n), \text{Ber}(p))$ between $\text{Ber}(\bar{X}_n)$ and $\text{Ber}(p)$ as a function of the random variable \bar{X}_n , and hence a random variable itself. Does $\text{TV}(\text{Ber}(\bar{X}_n), \text{Ber}(p))$ necessarily converge in probability to a constant? If yes, enter the constant below; if not; enter DNE.

$$\text{TV}(\text{Ber}(\bar{X}_n), \text{Ber}(p)) \xrightarrow[n \rightarrow \infty]{(\mathbf{P})} \boxed{0} \quad \checkmark$$

0

STANDARD NOTATION

Submit

You have used 1 of 2 attempts

✓ Correct (2/2 points)

(c)

1/1 point (graded)

Compute the total variation distance between

$$P = \text{Unif}([0, s]) \quad \text{and} \quad Q = \text{Unif}([0, t]), \quad \text{where } 0 < s < t.$$

$$\text{TV}(\mathbf{P}, \mathbf{Q}) = \boxed{1-s/t} \quad \checkmark$$

$1 - \frac{s}{t}$

STANDARD NOTATION

Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)

(d)

1/1 point (graded)

Let $X \sim N(\mu, \sigma^2)$ and $Y \sim \text{Ber}(p)$. Compute the total variation distance between the distributions of $\text{sign}(X)$ and $Y - 1$. Note that $\text{sign}(X)$ is a function of the random variable with

$$\text{sign}(X) = \begin{cases} 1 & \text{if } X > 0 \\ 0 & \text{if } X = 0 \\ -1 & \text{if } X < 0. \end{cases}$$

(If applicable, enter **abs(x)** for $|x|$, **Phi(x)** for $\Phi(x) = \mathbf{P}(Z \leq x)$ where $Z \sim \mathcal{N}(0, 1)$, and **q(alpha)** for q_α , the $1 - \alpha$ -quantile of a standard normal distribution, e.g. enter **q(0.01)** for $q_{0.01}$.)

$\text{TV}(\text{sign}(X), Y - 1) =$



STANDARD NOTATION

Submit

You have used 2 of 3 attempts

✓ Correct (1/1 point)

(e)

1/1 point (graded)

Compute the total variation distance between

$$\mathbf{P} = \text{Ber}(p) \quad \text{and} \quad \mathbf{Q} = \text{Pois}(p), \quad \text{where } p \in (0, 1).$$

TV(P, Q) =

$$p - p \cdot e^{-p}$$



$$p - p \cdot e^{-p}$$

STANDARD NOTATION

Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)

Discussion

Hide Discussion

Topic: Unit 3 Methods of Estimation: Homework 4: TV distance, KL-Divergence, and Introduction to MLE / 2.
Compute Total Variation Distance

Add a Post

[◀ All Posts](#)

Not getting why I got Part (d) not accepted

discussion posted 21 minutes ago by [sandipan dey](#).

I got my answer as $(1/4 + p/2) + |1/4 - p/2|$ for Part (d), but it's not accepted by the grader, any hint in terms of whether I am in the right direction? thank you very much in advance. It can also be written as $\max(1/2, p)$ but I think it will again get rejected by the grader.

[EDIT] my bad, i wrongly assumed $X \sim N(0, 1)$. Now it got accepted.

This post is visible to everyone.



0 responses

Preview

Submit

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

Learn About Verified Certificates

© All Rights Reserved