

Audit Access Expires Sep 9, 2020

You lose all access to this course, including your progress, on Sep 9, 2020.

Upgrade by Jul 5, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

A Proof of Cantor's Theorem

Let us refer to the set of A 's subsets as A 's **powerset**; in symbols: $\mathcal{P}(A)$. Cantor's Theorem can then be restated as follows:

Cantor's Theorem

For any set A , $|A| < |\mathcal{P}(A)|$

In order to prove this result, we'll need to verify each of the following two statements:

1. $|A| \leq |\mathcal{P}(A)|$
2. $|A| \neq |\mathcal{P}(A)|$

The first of these statements is straightforward, since the function $f(x) = \{x\}$ is an injection from A to $\mathcal{P}(A)$.

So what we really need to verify is the second statement: $|A| \neq |\mathcal{P}(A)|$.

Our proof of $|A| \neq |\mathcal{P}(A)|$ will proceed by *reductio*. We will assume the negation of what we wish to prove, and use this assumption to prove a contradiction. Since we want to prove $|A| \neq |\mathcal{P}(A)|$, this means that we will assume $|A| = |\mathcal{P}(A)|$. We will assume, in other words, that there is a bijection f from A to $\mathcal{P}(A)$.

Note that for each a in A , $f(a)$ is a member of $\mathcal{P}(A)$, and therefore a subset of A . So we can consider the question of whether a is a member of $f(a)$. In symbols:

$$a \in f(a)?$$

Note that this question will be answered negatively for whichever $a \in A$ is mapped by f to the empty set. Notice, moreover, that (if A is non-empty) the question will be answered positively for whichever member of A is mapped by f to A itself.

Let D be the set of members of A for which the question is answered negatively. In other words:

$$D = \{x \in A : x \notin f(x)\}$$

Since D is a subset of A , f must map some d in A to D . Now consider the following question:

$$d \in D?$$

The definition of D tells us that d is a member of D if and only if d is not a member of $f(d)$. In symbols:

$$d \in D \leftrightarrow d \notin f(d)$$


But the definition of d tells us that $f(d) = D$. So we have:

$$d \in D \leftrightarrow d \notin D$$

That statement is a contradiction: it tells us that something is the case if and only if it is not the case. So we have concluded our proof. We have assumed $|A| = |\mathcal{P}(A)|$, and used it to prove a contradiction. From this it follows that $|A| = |\mathcal{P}(A)|$ is false, and therefore that $|A| \neq |\mathcal{P}(A)|$ is true.

Cantor's Theorem is an amazing result. It shows that regardless of how many individuals there are in a set A , and regardless of whether A is finite or infinite A , there must be even more individuals in A 's power set. So there are infinitely many sizes of infinity!

Video Review: Proving Cantor's Theorem



▶ 0:00 / 0:00

▶ 1.50x 🔊 🗑️ 📄 🗣️

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Discussion

Topic: Week 1 / A Proof of Cantor's Theorem

Hide Discussion

Add a Post

Show all posts ▾

by recent activity ▾

✓ Empty set and its powerset

Would the cardinality of the empty set be equal to its power set in violation of Cantor's theorem? Or is the argument that: set A, the empty set, has a cardinality of 0 because it...

2 ▾

✓ Want more infinities

how to make even larger infinities because we can only apply the powerset a natural amount of times. what about when we can not do new method anymore

5 ▾

© All Rights Reserved