

< Previous





Next >

2.3.1 What is the Principle of Mathematical Induction?

 Bookmark this page

2.3.1 What is the Principle of Mathematical Induction?

The Principle of Mathematical Induction

©2014 R. van de Geijn and M. Myers

1 / 7

0:22 / 1:14

▶ 2.0x

🔊

🗒

📄

🗣

mathematical induction.

We're going to give you some practice proving various things via mathematical induction.

What's the principle of mathematical induction?

Well, there are actually several different ways of doing mathematical induction.

And what we're going to do is sometimes called weak induction.

So what do we have?

If one can show that some property holds for k equal to some base case.

In a previous example, this base case was equal to 1.

And we can show that if it holds for k equals K , where k is greater than or equal to that base case, then it also holds for the next little k .

Video
📄 [Download video file](#)

Transcripts
📄 [Download SubRip \(.srt\) file](#)
📄 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)
Read Unit 2.3.1 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

✓ Correct

Discussion

Topic: Week 2 / 2.3.1

Hide Discussion

Add a Post

◀ All Posts

Inductive set

question posted 2 years ago by [jchen9619](#)

What is the definition of a set that can be defined inductively?

+

⬆

🧮 Calculator

This post is visible to everyone.

Add a Response

1 response

maggiemyers (Staff)
about a year ago



From Cornell notes:

"An inductively defined set is a set where the elements are constructed by a finite number of applications of a given set of rules for creating more complicated objects from simpler ones."

My thinking:
Sets are collections. Some of theses sets can be "ordered" so that the entire collection can be described by rules (3 clauses.) They are to describe set S, specify

- one or more elements of S (basis clause). one or more rules to
- construct elements of S from existing elements of S (Inductive clause)
- that no other elements are in S (closure or extent clause).

While sets are collections where order doesn't matter, this ordering allows us to describe the contents of the set and then create arguments about it.

Add a comment

Showing all responses

Add a response:

Preview

Submit

Comments by Maggie in response to a posting on the discussion board

If mathematical induction intimidates you, have a look at in the enrichment (Unit 2.5.2) :Puzzles and Paradoxes in Mathematical Induction", by Adam Bjorndahl.

Here is my take on Induction. I am actually extending it beyond the proofs we do.

If you want to prove something holds for all members of a set that can be defined inductively, then you would use mathematical induction. You may recall a set is a collection and as such the order of its members is not important. However, some sets do have a natural ordering that can be used to describe the membership. This is especially valuable when the set has an infinite number of members, for example, natural numbers. Sets for which the membership can be described by suggesting there is a first element (or small group of firsts) then from this first you can create another (or others) then more and more by applying a rule to get another element in the set are our focus here. If all elements (members) are in the set because they are either the first (basis) or can be constructed by applying "The" rule to the first (basis) a finite number of times, then the set can be inductively defined.

So for us, the set of natural numbers is inductively defined. As a computer scientist you would say 0 is the first element and the rule is to add one to get another element. So 0,1,2,3,... are members of the natural numbers. 10 is a member of natural numbers.

numbers because you can find it by adding 1 to 0 ten times to get it.

So, the Principle of Mathematical induction proves that something is true for all of the members of a set that can be defined inductively. If this set has an infinite number of members, you couldn't show it is true for each of them individually. The idea is if it is true for the first(s) and it is true for any constructed member(s) no matter where you are in the list, it must be true for all. Why? Since we are proving things about natural numbers, the idea is if it is true for 0 and the next constructed, it must be true for 1 but then its true for 2, and then 3 and 4 and 5 ...and 10 and ...10000 and 10001, etc (all natural numbers) This is only because of the special ordering we can put on this set so we can know there is a next one for which it must be true. People often picture this rule by thinking of climbing a ladder or pushing down dominoes. If you know you started and you know where ever you are the next will follow then you must make it through all (even if there are an infinite number).

That is why to prove something using the Principle of Mathematical Induction you must show what you are proving holds at a start and then if it holds (assume it holds up to some point) then it holds for the next constructed element in the set. With these two parts shown, we know it must hold for all members of this inductively defined set.

You can find many examples of how to prove using PMI as well as many examples of when and why this method of proof will fail all over the web. Notice it only works for statements about sets "that can be defined inductively". Also notice subsets of natural numbers can often be defined inductively. For example, if I am a mathematician I may start counting at 1. Or I may decide that the statement holds for natural numbers ≥ 4 so I start my base case at 4.

< Previous

Next >



edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers
- News

Legal

- Terms of Service & Honor Code
- Privacy Policy
- Accessibility Policy
- Trademark Policy
- Sitemap
- Cookie Policy
- Your Privacy Choices

Connect

- Idea Hub
- Contact Us
- Help Center
- Security

Calculator

[Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)