



MITx 6.419x

Data Analysis: Statistical Modeling and Computation in Applications

Help

sandipan\_dey ▾

- [Course](#)
- [Progress](#)
- [Dates](#)
- [Discussion](#)
- [Resources](#)

[🏠 Course](#) / [Module 5: Environmental Data and Gaussian Processes](#) / [Spatial Prediction](#)



< Previous	✓	✓	✓	✓		✓	Next >
------------	---	---	---	---	--	---	--------

5. Effects of Parameters on Kernel Functions

Bookmark this page



between points, eg.  $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$  and  $\mathbf{x}' = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ , and  $r$  is a distance, eg.  $r = \|\mathbf{x} - \mathbf{x}'\|$ .

constant	$\sigma_0^2$
linear	$\sum_{d=1}^D \sigma_d^2 x_d x'_d$
polynomial	$(\mathbf{x} \cdot \mathbf{x}' + \sigma_0^2)^p$
squared exponential	$\exp\left(-\frac{r^2}{2\ell^2}\right)$
Matérn	$\frac{1}{2^{\nu-1}\Gamma(\nu)}\left(\frac{\sqrt{2\nu}}{\ell}r\right)^\nu K_\nu\left(\frac{\sqrt{2\nu}}{\ell}r\right)$
exponential	$\exp\left(-\frac{r}{\ell}\right)$
$\gamma$ -exponential	$\exp\left(-\left[\frac{r}{\ell}\right]^\gamma\right)$
rational quadratic	$\left(1 + \frac{r^2}{2\alpha\ell^2}\right)^{-\alpha}$
neural network	$\sin^{-1}\left(\frac{2\mathbf{x}^\top\boldsymbol{\Sigma}\mathbf{x}}{\sqrt{(1+2\mathbf{x}^\top\boldsymbol{\Sigma}\mathbf{x})(1+2\mathbf{x}'^\top\boldsymbol{\Sigma}\mathbf{x}')}}\right)$


Note, for example, that the last kernel in this table is stationary, but it is not isotropic as the matrix  $\boldsymbol{\Sigma}$  means that the kernel is no longer a function of purely distance. Rather, the distance between two points is scaled depending on the direction of the vector between them; hence, the kernel is not isotropic.

For example, consider the following covariance function:

$$k(Z_1, Z_2) = \exp\left(-\left[\frac{\|Z_1 - Z_2\|}{\ell}\right]^\gamma\right).$$

This is the  $\gamma$ -exponential shown in table above.

Examples of this function are shown below for various choices of  $\gamma$ .



**34:** The effects of the parameter  $\gamma$  on the covariance function.

When  $\gamma = 1$ , the peak of the function is sharp, and the function is heavy tailed. For  $\gamma = 2$  the function is smooth around the peak, but falls off sharply at large distance. Although  $\gamma = 2$  is a common choice, we can make other choices if we have reason to suspect that the observations are heavily correlated at large distance.

We can also combine kernel functions to form new kernel functions.

If  $k_1$  is a kernel function, and  $k_2$  is also a kernel function, then

$$k(Z_1, Z_2) = k_1(Z_1, Z_2) + k_2(Z_1, Z_2)$$

is also a kernel function.

Similarly,

$$k(Z_1, Z_2) = k_1(Z_1, Z_2) \times k_2(Z_1, Z_2)$$

is also a kernel function.

### Combinations of kernel functions 1

1/1 point (graded)  
Suppose that  $k_1$  and  $k_2$  are both isotropic kernel functions.

Is the linear combination

$$k(Z_1, Z_2) = 2k_1(Z_1, Z_2) + k_2(Z_1, Z_2)$$

also an isotropic kernel function?

☒ Yes

☐ No



**Solution:**

If  $k_1$  and  $k_2$  are isotropic, then they only depend on the distance between the supplied points. Therefore  $k$  also only depends on this distance, and nothing else. Thus  $k$  is also isotropic.

Submit

You have used 1 of 1 attempt

Answers are displayed within the problem

### Combinations of kernel functions 2

1/1 point (graded)  
Suppose that  $k_1$  is an isotropic kernel function, but  $k_2$  is a stationary kernel function.

What can we say about the product

$$k(Z_1, Z_2) = k_1(Z_1, Z_2) \times k_2(Z_1, Z_2)$$

It is

☒ Stationary

☐ Isotropic

☐ None of the above



**Solution:**

Although all isotropic kernel functions must be stationary, not all stationary kernel functions are isotropic. Thus, we can only say that a combination of a stationary and isotropic kernel will be stationary.

Submit

You have used 1 of 2 attempts

**i** Answers are displayed within the problem

For a comprehensive study of covariance functions and kernels, the reader should check Chapter 4 in Williams, Christopher KI, and Carl Edward Rasmussen. Gaussian processes for machine learning. Vol. 2. No. 3. Cambridge, MA: MIT Press, 2006.

Discussion

Hide Discussion

Topic: Module 5: Environmental Data and Gaussian Processes:Spatial Prediction / 5.  
Effects of Parameters on Kernel Functions

Add a Post

Show all posts	by recent activity
<div>Too difficult compared to video</div> <div>I feel like the theoretical question are way too difficult compared to the video courses.</div> <div>1</div>	
<div>Is the length parameter constant?</div> <div>In the definition of RBF kernel, could the length parameter (l) be variable depending on x and x' ? If we apply GP with graph network ...</div> <div>1</div>	

< Previous

Next >



## edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers
- News

## Legal

- Terms of Service & Honor Code
- Privacy Policy
- Accessibility Policy
- Trademark Policy
- Sitemap

# Connect

[Blog](#)

[Contact Us](#)

[Help Center](#)

[Media Kit](#)

[Donate](#)



© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)