EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the <u>Privacy Policy</u>.





Homework 3: Introduction to

4. Relating Hypothesis Tests and

Course > Unit 2 Foundation of Inference > Hypothesis Testing

> Confidence intervals

Currently enrolled in **Audit Track** (expires December 25, 2019) <u>Upgrade (\$300)</u>

4. Relating Hypothesis Tests and Confidence intervals

(a)

2.0/2 points (graded)

Consider an i.i.d. sample $X_1,\ldots,X_n\sim \mathsf{Poiss}\,(\lambda)$ for $\lambda>0$.

Starting from the Central Limit Theorem, find a confidence interval I=[A,B] with asymptotic level $1-\alpha$ that is centered about \overline{X}_n using the plug-in method.

Write $\operatorname{barX_n}$ for \overline{X}_n . If applicable, type $\operatorname{abs(x)}$ for |x|, $\operatorname{Phi(x)}$ for $\Phi(x) = \mathbf{P}(Z \leq x)$ where $Z \sim \mathcal{N}(0,1)$, and $\operatorname{q(alpha)}$ for q_{α} , the $1-\alpha$ quantile of a standard normal variable.)

$$\mathcal{I} = [A,B]$$
 for

$$A=egin{array}{c} \mathsf{barX_n} \cdot \mathsf{q(alpha/2)*sqrt(barX_n/n)} \end{array}$$

$$B = barX_n + q(alpha/2)*sqrt(barX_n/n)$$

STANDARD NOTATION

Submit

You have used 1 of 3 attempts

(b)

2/2 points (graded)

Consider the following hypothesis with a fixed number $\,\lambda_0>0$:

$$H_0: \lambda = \lambda_0 \quad ext{vs} \quad H_1: \lambda
eq \lambda_0.$$

Define a test for the above hypotheses with asymptotic level α , and rewrite it in the form

$$\psi = \mathbf{1}\{\lambda_0
otin J\},$$

for some interval $\, J$.

$$\mathcal{J} = [C,D]$$
 for

$$C =$$
 barX_n - q(alpha/2)*sqrt(barX_n /n)

$$D = barX_n + q(alpha/2)*sqrt(barX_n /n)$$

Submit

You have used 1 of 3 attempts

✓ Correct (2/2 points)

Discussion

Topic: Unit 2 Foundation of Inference:Homework 3: Introduction to Hypothesis Testing / 4. Relating Hypothesis

Tests and Confidence intervals **Hide Discussion** Add a Post **≺** All Posts Question B: Using the solve method + discussion posted 2 days ago by sean s wang Basically I followed the process in recitation video #2. The professor did not finish the calculation. So I am just solving a quadratic equation to get the result. It's a little messy but I think it is right. I was a little careless and lost two attempts. Hence I want to double check to see if I am missing anything before my final submission. Thanks! This post is visible to everyone. 1 response Add a Response markweitzman (Community TA) 2 days ago The guestion asks for you to use plug-in method - much easier. "...using the plug-in method." ••• You are right. I used the wrong terminology. It should be plug-in method (although question B does not specifically say it). I should be able to just use the result of the previous question. posted a day ago by sean s wang ••• The reason I got it wrong is somehow I keep forgetting the square root operation for the the variance λ . Guess I just need to practice more. posted a day ago by sean s wang

It wasn't immediately obvious to me either that we weren't	supposed to use a different method for part b). I also had a huge me	ssy quadratic.
posted a day ago by synnfusion	The second of th	
Add a comment		
. "		
owing all responses		
dd a response:		
eview		
Submit		
dit Assocs Evnivos Dos 24, 2010		
dit Access Expires Dec 24, 2019 u lose all access to this course, including your		
grade by Nov 4, 2019 to get unlimited access	to the course as long as it exists on the site. <u>Upgrad</u>	de now
	Learn About Verified Certificates	
		© All Rights Re