



< Previous



Next >

3. Matrices

Bookmark this page



Calculator



Hide Notes

Problem Set B due Sep 15, 2021 20:30 IST



Practice

Rotation determinant

1/1 point (graded)
Let R_θ be the matrix that rotates a two-dimensional vector by θ counter-clockwise (that is, the vector $R_\theta \vec{v}$ is obtained by rotating \vec{v} counter-clockwise by the angle θ). Find the determinant of R_θ .

$\det R_\theta =$ ✓ Answer: 1

Solution:

We have

$$R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \tag{5.222}$$

Therefore the determinant is given by $\cos^2 \theta + \sin^2 \theta$, which equals **1**.

Submit

You have used 1 of 3 attempts

ⓘ Answers are displayed within the problem

Inverse Rotation

1.0/1 point (graded)
Find the inverse of the rotation matrix R_θ . Type for θ .

(Enter a matrix using notation such as .)

✓

? INPUT HELP

Submit

You have used 1 of 3 attempts

Matrix Transpose

1.0/1 point (graded)
Given a matrix A , we write A^T for the matrix whose entry in row i and column j is given by the entry in row j and column i of A . This A^T is called the *transpose* of A . For example,

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \iff A^T = \begin{pmatrix} 1 & 3 \\ 2 & 4 \end{pmatrix} \tag{5.223}$$

Visually, one obtains the transpose by “reflecting” the matrix entries across the top-left to bottom-right diagonal.

Let $M = \begin{pmatrix} 0 & 1 \\ 2 & -1 \end{pmatrix}$. Find M^T .

(Enter a matrix using notation such as `[[a,b],[c,d]]`.)

$M^T =$

`[[0,2],[1,-1]]`

✓ Answer: `[[0, 2],[1,-1]]`

Solution:

For a 2×2 matrix, we interchange the lower-left and top-right entries to obtain the transpose $\begin{pmatrix} 0 & 2 \\ 1 & -1 \end{pmatrix}$.

Submit

You have used 1 of 3 attempts

Answers are displayed within the problem

Orthogonal Matrices

1/1 point (graded)
A square matrix A is called *orthogonal* if $A^T = A^{-1}$. Which of the following matrices are orthogonal? Choose all that apply.

- ☐ $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$
- ☒ R_θ (rotation matrix)
- ☒ $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$
- ☒ $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$



Solution:

The answer is found by computing A^T and A^{-1} for each matrix A and seeing if they are equal.

Submit

You have used 1 of 3 attempts

Answers are displayed within the problem

Find Orthogonal Matrices

4/4 points (graded)
Find four distinct orthogonal 2×2 matrices, each of which has top-left entry equal to $\frac{-1}{\sqrt{2}}$.

(Enter a matrix using notation such as `[[a,b],[c,d]]`.)

- `[[-1/sqrt(2), -1/sqrt(2)], [1/sqrt(2), -1/sqrt(2)]]`
- ✓ Answer: `[[-1/sqrt(2) , -1/sqrt(2)], [1/sqrt(2) , -1/sqrt(2)]]`
- `[[-1/sqrt(2), 1/sqrt(2)], [-1/sqrt(2), -1/sqrt(2)]]`
- ✓ Answer: `[[-1/sqrt(2) , 1/sqrt(2)], [-1/sqrt(2) , -1/sqrt(2)]]`
- `[[-1/sqrt(2), 1/sqrt(2)], [1/sqrt(2), 1/sqrt(2)]]`
- ✓ Answer: `[[-1/sqrt(2) , 1/sqrt(2)], [1/sqrt(2) , 1/sqrt(2)]]`

[[-1/sqrt(2), 1/sqrt(2)], [1/sqrt(2), 1/sqrt(2)]]

[[-1/sqrt(2), -1/sqrt(2)], [-1/sqrt(2), 1/sqrt(2)]]

▼ **Answer:** [[-1/sqrt(2) , 1/sqrt(2)], [1/sqrt(2) , 1/sqrt(2)]]

✓ **Answer:** [[-1/sqrt(2) , -1/sqrt(2)], [-1/sqrt(2) , 1/sqrt(2)]]

Solution:

For a 2×2 matrix to be orthogonal, we need $A^{-1} = A^T$, in symbols:

$$\begin{pmatrix} a & c \\ b & d \end{pmatrix} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

(5.224)

By matching the top-left or bottom-right entries, we see that $ad - bc$ equals either 1 (case 1) or -1 (case 2).

In case 1, we have $d = a$ and $c = -b$, so the $ad - bc = 1$ constraint becomes $a^2 + b^2 = 1$. With $a = \frac{-1}{\sqrt{2}}$, the solutions are $b = \pm \frac{1}{\sqrt{2}}$.

$$M_0 = \begin{pmatrix} \frac{-1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \end{pmatrix} \quad \text{or} \quad M_1 = \begin{pmatrix} \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{-1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \end{pmatrix}$$

(5.225)

In fact, these are the rotation matrices $R^{3\pi/4}$ and $R^{-3\pi/4}$.

In case 2, we have $d = -a$ and $c = b$, so the $ad - bc = -1$ constraint becomes $-a^2 - b^2 = -1$. With $a = \frac{-1}{\sqrt{2}}$, again we get the solutions $b = \pm \frac{1}{\sqrt{2}}$. The corresponding matrices are:

$$M_2 = \begin{pmatrix} \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \quad \text{or} \quad M_3 = \begin{pmatrix} \frac{-1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$$

(5.226)

It follows that these are the only four orthogonal 2×2 matrices with $a = \frac{-1}{\sqrt{2}}$.

Submit

You have used 1 of 3 attempts

📘 Answers are displayed within the problem

3. Matrices

Hide Discussion

Topic: Unit 4: Matrices and Linearization / 3. Matrices

Add a Post

Show all posts ▼by recent activity ▼

🗨️ [STAFF] Could not format HTML for problem. Contact course staff in the discussion forum for assistance.

For the 2nd and 3rd problem, it shows "Could not format HTML for problem. Contact course staff in the discussion forum for assista... 1 new_

★ Following



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)





© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)