EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the <u>Privacy Policy</u>.

×



<u>Lecture 10: Consistency of MLE,</u>

Covariance Matrices, and

Course > Unit 3 Methods of Estimation > Multivariate Statistics

> 5. Review: Covariance

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

5. Review: Covariance

Note: The following exercises are a review of covariance, and will be discussed in lecture. We encourage that you attempt these exercises before watching the video.

Review: Covariance

2/2 points (graded)

If X and Y are random variables with respective means μ_X and μ_Y , then recall the **covariance** of X and Y (written $\mathsf{Cov}\,(X,Y)$) is defined to be

$$\mathsf{Cov}\left(X,Y
ight) = \mathbb{E}\left[\left(X - \mu_X
ight)\left(Y - \mu_Y
ight)
ight].$$

Alternatively, one can show that this is equivalent to $\mathbb{E}\left[XY\right] - \mathbb{E}\left[X\right]\mathbb{E}\left[Y\right]$.

For each of the following statements, indicate whether it is true or false.

 $"\mathsf{Cov}\,(X,X)=\mathsf{Var}\,(X)".$

True

False



"Like the variance, the covariance between an arbitrary pair of RVs X and Y is always non-negative."







Solution:

- True. Cov $(X,X)=\mathbb{E}\left[(X-\mu_X)^2\right]=\mathsf{Var}\left(X\right)$.
- False. Consider (X,Y) which is distributed uniformly over the set $\{(1,-1),(-1,1)\}$. The marginal distributions of both X and Y are uniform over $\{\pm 1\}$, so $\mu_X=\mu_Y=0$. On the other hand, $\mathbb{E}\left[XY\right]=-1$, so $\mathsf{Cov}\left(X,Y\right)=-1$.

Submit

You have used 1 of 1 attempt

• Answers are displayed within the problem

Alternate Formula for Covariance

0/1 point (graded)

Let X and Y are random variables with respective means μ_X and μ_Y . Is it true that $\mathbb{E}\left[(X)\left(Y-\mu_Y
ight)
ight]=\mathsf{Cov}\left(X,Y
ight)$?







Solution:

Indeed, $\mathbb{E}\left[\left(X\right)\left(Y-\mu_Y\right)\right]=\mathsf{Cov}\left(X,Y\right)=\mathbb{E}\left[\left(X-\mu_X\right)\left(Y\right)\right]$. That is, it is sufficient to center one random variable around its mean when computing the covariance between two random variables. This can be seen from the following:


$$\mathbb{E}\left[\left(X\right)\left(Y-\mu_{Y}\right)\right] = \mathbb{E}\left[XY\right] - \mathbb{E}\left[X\mu_{Y}\right]$$
$$= \mathbb{E}\left[XY\right] - \mathbb{E}\left[X\right]\mathbb{E}\left[Y\right].$$

Submit

You have used 1 of 1 attempt

1 Answers are displayed within the problem

Covariance: Definition and Formula



Video

<u>Download video file</u>

Transcripts

<u>Download SubRip (.srt) file</u>

<u>Download Text (.txt) file</u>


Bilinearity of Covariance

1/1 point (graded)

Let X, Y, Z be random variables and a, b be constants. Indicate whether the following statement is true or false.

"Covariance is bilinear, i.e. $\mathsf{Cov}\left(aX+bY,Z\right)=a\mathsf{Cov}\left(X,Z\right)+b\mathsf{Cov}\left(Y,Z\right)$."

Hint: Use the result from the problem immediately above.



✓ Correct (1/1 point)

Example of Covariance I

1/1 point (graded)

Let A=X+Y and B=X-Y. Let $\mu_X=\mathbb{E}\left[X\right]$, $\mu_Y=\mathbb{E}\left[Y\right]$, $\tau_X=\mathsf{Var}\left(X\right)$, $\tau_Y=\mathsf{Var}\left(Y\right)$ and $c=\mathsf{Cov}\left(X,Y\right)$. In terms of $\mu_X,\,\mu_Y,\,\tau_X,\,\tau_Y,\,$ and $c,\,$ what is $\mathsf{Cov}\left(A,B\right)$?

(Enter **mu_X** for μ_X , **tau_X** for τ_X .)

tau_X-tau_Y
$$\checkmark$$
 Answer: tau_X-tau_Y
$$\tau_X - \tau_Y$$
 STANDARD NOTATION

Solution:

Expand out the definition of covariance using bi-linearity (see the solution to the previous question):

$$\begin{aligned} \mathsf{Cov}\left(A,B\right) &= \mathsf{Cov}\left(X+Y,X-Y\right) \\ &= \mathsf{Cov}\left(X+Y,X\right) - \mathsf{Cov}\left(X+Y,Y\right) \\ &= \mathsf{Cov}\left(X,X\right) + \mathsf{Cov}\left(Y,X\right) - \mathsf{Cov}\left(X,Y\right) - \mathsf{Cov}\left(Y,Y\right) \\ &= \mathsf{Var}\left(X\right) - \mathsf{Var}\left(Y\right) \\ &= \tau_X - \tau_Y. \end{aligned}$$

Submit

You have used 1 of 4 attempts

• Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 3 Methods of Estimation:Lecture 10: Consistency of MLE, Covariance Matrices, and Multivariate Statistics / 5. Review: Covariance

Add a Post

Show all posts ▼

All terms required in 2nd question?

possible error in first question?

If my browser is correct, what I see is that the question is Cov(x,x) = Var(x)^n. I see an "n". and based on that I answered ... although the grader is marking me a red although d...

The possible error in first question?

If my browser is correct, what I see is that the question is Cov(x,x) = Var(x)^n. I see an "n". and based on that I answered ... although the grader is marking me a red although d...

Learn About Verified Certificates

© All Rights Reserved