

MITx: 14.310x Data Analysis for Social Scientists

Heli



Bookmarks

- Module 1: The Basics of R and Introduction to the Course
- ► Entrance Survey
- Module 2: Fundamentals of Probability, Random Variables, Distributions, and Joint Distributions
- Module 3: Gathering and Collecting Data,
 Ethics, and Kernel
 Density Estimates
- Module 4: Joint,
 Marginal, and
 Conditional
 Distributions &
 Functions of Random
 Variable

Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation > Human Subjects and Special Distributions > The Binomial Distribution - Quiz

The Binomial Distribution - Quiz

☐ Bookmark this page

Question 1

1/1 point (graded)

Which of the following is the appropriate interpretation of the Binomial distribution?

- ullet a. The distribution of a series of trials, each with any number of outcomes $1,\ldots,k$ and associated probabilities p_1,\ldots,p_k
- ullet b. The number of successes in a sequence of n success/failure trials, each of which may have a different associated probability, p_1, \ldots, p_n
- ullet c. The distribution of the different probabilities of success, p_1,\ldots,p_n for a series of n success/failure trials
- \odot d. The number of successes in a sequence of n success/failure trials, each of which has the same probability of success, p

Explanation

- Module 5: Moments of a Random Variable,
 Applications to Auctions,
 Intro to Regression
- Module 6: Special
 <u>Distributions, the</u>

 <u>Sample Mean, the</u>
 <u>Central Limit Theorem,</u>
 and Estimation

<u>Human Subjects and</u> <u>Special Distributions</u>

Finger Exercises due Nov 07, 2016 at 05:00 IST

The Sample Mean, Central Limit Theorem, and Estimation

Finger Exercises due Nov 07, 2016 at 05:00 IST

Module 6: Homework

<u>Homework due Oct 31, 2016 at</u> 05:00 IST

(A)

▶ Exit Survey

A binomial distribution refers the number of successes in a sequence of n success/failure trials (meaning, there are only two outcomes) where the probability of success is constant across trials and given by p.

Submit

You have used 1 of 2 attempts

Correct (1/1 point)

Question 2

1/1 point (graded)

Let's go back to the example of Anna and Brian as soccer players. As before, the probability that Anna makes a shot is $p_A=0.2$ and the probability that Brian makes a shot is $p_B=0.5$.

Suppose that Anna takes 50 shots and that Brian takes 25 shots. For each player, the probability that they will make a shot does not change over time. In other words, each of their 25 to 50 shots are independent.


Calculate the expectation and variance for each of Anna's and Brian's shots according to the sequence of shots attempted as described above.

Provide exact answers, i.e. do not round

Expectation of Anna's 50 shots


10

✓ Answer: 10



Explanation

The expectation of Anna's 50 shots is given by np=50*0.2=10 and the variance is given by npq=50*0.2*0.8=8. The expectation of Brian's 25 shots is given by np=25*0.5=12.5 and the variance is given by npq=25*0.5*0.5=6.25.



© All Rights Reserved



© 2016 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.















