

This website uses cookies to ensure you get the best experience on our website.

Ok [Learn more](#)

# STHDA

Statistical tools for high-throughput data analysis

Licence:

[Home](#)[Basics](#)[Data](#)[Visualize](#)[Analyze](#)[Products](#)[Contribute](#)[Support](#)[About](#)

## Sign in

### Login

### Password

Auto connect ☒

Register

Forgotten password

## Welcome!

Want to Learn More on R  
Programming and Data  
Science?

Follow us [by Email](#)

by [FeedBurner](#)

on Social Networks

[Home](#) / [Easy Guides](#) / [R software](#) / [R Basic Statistics](#) / [Comparing Means in R](#) / [Unpaired Two-Samples T-test in R](#)



## Book your next getaway


Escape to Bali on your  
next adventure with your  
ICIC card

Malaysia Airlines

## Unpaired Two-Samples T-test in R



- [What is unpaired two-samples t-test?](#)
- [Research questions and statistical hypotheses](#)
- [Formula of unpaired two-samples t-test](#)
- [Visualize your data and compute unpaired two-samples t-test in R](#)
  - [Install ggpubr R package for data visualization](#)
  - [R function to compute unpaired two-samples t-test](#)
  - [Import your data into R](#)
  - [Check your data](#)
  - [Visualize your data using box plots](#)
  - [Preliminary test to check independent t-test assumptions](#)
  - [Compute unpaired two-samples t-test](#)

 Ads by Google[analyzing data](#)[r unpaired t test](#)[two tailed t test](#)[free test exams](#)[11 sample tests](#)

### R Packages

 [factoextra](#) [survminer](#) [ggpubr](#) [ggcorrplot](#) [fastqcr](#)

- [Interpretation of the result](#)
- [Access to the values returned by t.test\(\) function](#)

- [Online unpaired two-samples t-test calculator](#)
- [See also](#)
- [Infos](#)

## What is unpaired two-samples t-test?

The **unpaired two-samples t-test** is used to compare the **mean** of two independent groups.

For example, suppose that we have measured the weight of 100 individuals: 50 women (group A) and 50 men (group B). We want to know if the mean weight of women ( $m_A$ ) is significantly different from that of men ( $m_B$ ).




In this case, we have two unrelated (i.e., independent or unpaired) groups of samples. Therefore, it's possible to use an **independent t-test** to evaluate whether the means are different.



Note that, unpaired two-samples t-test can be used only under certain conditions:

- when the two groups of samples (A and B), being compared, are **normally distributed**. This can be checked using **Shapiro-Wilk test**.
- and when the **variances** of the two groups are equal. This can be checked using **F-test**.






## Our Books

### 3D Plots in R

## Unpaired Two-Samples T-test in R

- + Definition
- + Research Questions & Statistics
- + T-test Formula
- + Practical Examples in R
- + Interpret



© sthda.com 2016



This article describes the formula of the independent t-test and provides practical examples in R.

## Research questions and statistical hypotheses

Typical research questions are:




1. whether the mean of group A ( $m_A$ ) is equal to the mean of group B ( $m_B$ )?
2. whether the mean of group A ( $m_A$ ) is less than the mean of group B ( $m_B$ )?
3. whether the mean of group A ( $m_A$ ) is greater than the mean of group B ( $m_B$ )?

In statistics, we can define the corresponding *null hypothesis* ( $H_0$ ) as follow:

1.  $H_0 : m_A = m_B$
2.  $H_0 : m_A \leq m_B$
3.  $H_0 : m_A \geq m_B$

The corresponding *alternative hypotheses* ( $H_a$ ) are as follow:


1.  $H_a : m_A \neq m_B$  (different)
2.  $H_a : m_A > m_B$  (greater)



R Graphics Essentials for  
Great Data Visualization:  
200 Practical Examples  
You Want to Know for  
Data Science  
★ **NEW!!**



Practical Guide to Cluster  
Analysis in R



Practical Guide to  
Principal Component  
Methods in R

3.  $H_a : m_A < m_B$  (less)

! Note that:

- Hypotheses 1) are called **two-tailed tests**
- Hypotheses 2) and 3) are called **one-tailed tests**

## Formula of unpaired two-samples t-test

### 1. Classical t-test:

! If the variance of the two groups are equivalent (**homoscedasticity**), the **t-test value**, comparing the two samples ( $A$  and  $B$ ), can be calculated as follow.

$$t = \frac{m_A - m_B}{\sqrt{\frac{S^2}{n_A} + \frac{S^2}{n_B}}}$$

where,

- $m_A$  and  $m_B$  represent the mean value of the group A and B, respectively.
- $n_A$  and  $n_B$  represent the sizes of the group A and B, respectively.
- $S^2$  is an estimator of the pooled **variance** of the two groups. It can be calculated as follow :

$$S^2 = \frac{\sum (x - m_A)^2 + \sum (x - m_B)^2}{n_A + n_B - 2}$$

with degrees of freedom (df):  $df = n_A + n_B - 2$ .

### 2. Welch t-statistic:

! If the variances of the two groups being compared are different (**heteroscedasticity**), it's possible to use the **Welch t test**, an adaptation of Student t-test.


**Welch t-statistic** is calculated as follow :

$$t = \frac{m_A - m_B}{\sqrt{\frac{S_A^2}{n_A} + \frac{S_B^2}{n_B}}}$$

where,  $S_A$  and  $S_B$  are the standard deviation of the the two groups A and B, respectively.

! Unlike the classic Student's t-test, **Welch t-test formula** involves the variance of each of the two groups ( $S_A^2$  and  $S_B^2$ ) being compared. In other words, it does not use the pooled variance  $S$ .

The **degrees of freedom** of **Welch t-test** is estimated as follow :



### Guest Book

great

By Visitor

[Guest Book](#)

$$df = \left( \frac{S_A^2}{n_A} + \frac{S_B^2}{n_B} \right) / \left( \frac{S_A^4}{n_A^2(n_B - 1)} + \frac{S_B^4}{n_B^2(n_A - 1)} \right)$$



A p-value can be computed for the corresponding absolute value of t-statistic ( $|t|$ ).



Note that, the Welch t-test is considered as the safer one. Usually, the results of the **classical t-test** and the **Welch t-test** are very similar unless both the group sizes and the standard deviations are very different.



How to interpret the results?



If the p-value is inferior or equal to the significance level 0.05, we can reject the null hypothesis and accept the alternative hypothesis. In other words, we can conclude that the mean values of group A and B are significantly different.

## Visualize your data and compute unpaired two-samples t-test in R

### Install ggpubr R package for data visualization

You can draw R base graphs as described at this link: [R base graphs](#). Here, we'll use the **ggpubr R package** for an easy ggplot2-based data visualization

- Install the latest version from GitHub as follow (recommended):

```
# Install
if(!require(devtools)) install.packages("devtools")
devtools::install_github("kassambara/ggpubr")
```

- Or, install from CRAN as follow:

```
install.packages("ggpubr")
```

### R function to compute unpaired two-samples t-test

To perform two-samples t-test comparing the means of two independent samples (x & y), the R function **t.test()** can be used as follow:



DOES YOUR  
BUSINESS  
ACCEPT  
PAYPAL?



Blogroll

 **Datanovia: Online Data Science Courses**

 **R-Bloggers**

```
t.test(x, y, alternative = "two.sided", var.equal = FALSE)
```

- **x,y:** numeric vectors
- **alternative:** the alternative hypothesis. Allowed value is one of "two.sided" (default), "greater" or "less".
- **var.equal:** a logical variable indicating whether to treat the two variances as being equal. If TRUE then the pooled variance is used to estimate the variance otherwise the Welch test is used.

## Import your data into R

1. **Prepare your data** as specified here: [Best practices for preparing your data set for R](#)
2. **Save your data** in an external .txt tab or .csv files
3. **Import your data into R** as follow:

```
# If .txt tab file, use this
my_data <- read.delim(file.choose())
# Or, if .csv file, use this
my_data <- read.csv(file.choose())
```

Here, we'll use an example data set, which contains the weight of 18 individuals (9 women and 9 men):

```
# Data in two numeric vectors
women_weight <- c(38.9, 61.2, 73.3, 21.8, 63.4, 64.6, 48.4, 48.8, 48.5)
men_weight <- c(67.8, 60, 63.4, 76, 89.4, 73.3, 67.3, 61.3, 62.4)
# Create a data frame
my_data <- data.frame(
  group = rep(c("Woman", "Man"), each = 9),
  weight = c(women_weight, men_weight)
)
```

? We want to know, if the average women's weight differs from the average men's weight?

## Check your data

```
# Print all data
print(my_data)
```

```
  group weight
1  Woman   38.9
2  Woman   61.2
3  Woman   73.3
4  Woman   21.8
5  Woman   63.4
6  Woman   64.6
7  Woman   48.4
8  Woman   48.8
9  Woman   48.5
10 Man    67.8
11 Man    60.0
12 Man    63.4
13 Man    76.0
14 Man    89.4
15 Man    73.3
16 Man    67.3
17 Man    61.3
18 Man    62.4
```



It's possible to compute summary statistics (mean and sd) by groups. The dplyr package can be used.

- To install **dplyr** package, type this:

```
install.packages("dplyr")
```

- Compute summary statistics by groups:


```
library(dplyr)
group_by(my_data, group) %>%
  summarise(
    count = n(),
    mean = mean(weight, na.rm = TRUE),
    sd = sd(weight, na.rm = TRUE)
  )
```

```
Source: local data frame [2 x 4]
  group count    mean    sd
  (fctr) (int)  (dbl)  (dbl)
1  Man     9 68.98889 9.375426
2  Woman   9 52.10000 15.596714
```



## Visualize your data using box plots

```
# Plot weight by group and color by group
library("ggpubr")
ggboxplot(my_data, x = "group", y = "weight",
          color = "group", palette = c("#00AFBB", "#E7B800"),
          ylab = "Weight", xlab = "Groups")
```



## Preliminary test to check independent t-test assumptions

? **Assumption 1:** Are the two samples independent?

Yes, since the samples from men and women are not related.

? **Assumption 2:** Are the data from each of the 2 groups follow a normal distribution?

Use Shapiro-Wilk normality test as described at: [Normality Test in R](#). - Null hypothesis: the data are normally distributed - Alternative hypothesis: the data are not normally distributed

We'll use the functions **with()** and **shapiro.test()** to compute Shapiro-Wilk test for each group of samples.



```
# Shapiro-Wilk normality test for Men's weights
with(my_data, shapiro.test(weight[group == "Man"]))# p = 0.1
# Shapiro-Wilk normality test for Women's weights
with(my_data, shapiro.test(weight[group == "Woman"])) # p = 0.6
```



From the output, the two p-values are greater than the significance level 0.05 implying that the distribution of the data are not significantly different from the normal distribution. In other words, we can assume the normality.



Note that, if the data are not normally distributed, it's recommended to use the non parametric two-samples Wilcoxon rank test.



**Assumption 3.** Do the two populations have the same variances?

We'll use **F-test** to test for homogeneity in variances. This can be performed with the function **var.test()** as follow:

```
res.ftest <- var.test(weight ~ group, data = my_data)
res.ftest
```

```
F test to compare two variances
data: weight by group
F = 0.36134, num df = 8, denom df = 8, p-value = 0.1714
alternative hypothesis: true ratio of variances is not equal to 1
95 percent confidence interval:
 0.08150656 1.60191315
sample estimates:
ratio of variances
 0.3613398
```



The p-value of **F-test** is  $p = 0.1713596$ . It's greater than the significance level  $\alpha = 0.05$ . In conclusion, there is no significant difference between the variances of the two sets of data. Therefore, we can use the classic **t-test** with assume equality of **the two variances**.

## Compute unpaired two-samples t-test



Question : Is there any significant difference between women and men weights?

**1) Compute independent t-test - Method 1:** The data are saved in two different numeric vectors.

```
# Compute t-test
res <- t.test(women_weight, men_weight, var.equal = TRUE)
res
```

```
Two Sample t-test
data: women_weight and men_weight
t = -2.7842, df = 16, p-value = 0.01327
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 -29.748019 -4.029759
sample estimates:
mean of x mean of y
 52.10000  68.98889
```

**2) Compute independent t-test - Method 2:** The data are saved in a data frame.

```
# Compute t-test
res <- t.test(weight ~ group, data = my_data, var.equal = TRUE)
res
```

```
Two Sample t-test
data: weight by group
t = 2.7842, df = 16, p-value = 0.01327
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 4.029759 29.748019
sample estimates:
mean in group Man mean in group Woman
 68.98889          52.10000
```

! As you can see, the two methods give the same results.

In the result above :

- **t** is the **t-test statistic** value (t = 2.784),
- **df** is the degrees of freedom (df= 16),
- **p-value** is the significance level of the **t-test** (p-value = 0.01327).
- **conf.int** is the **confidence interval** of the mean at 95% (conf.int = [4.0298, 29.748]);
- **sample estimates** is the mean value of the sample (mean = 68.988889, 52.1).



Note that:

- if you want to test whether the average men's weight is less than the average women's weight, type this:

```
t.test(weight ~ group, data = my_data,  
       var.equal = TRUE, alternative = "less")
```

- Or, if you want to test whether the average men's weight is greater than the average women's weight, type this

```
t.test(weight ~ group, data = my_data,  
       var.equal = TRUE, alternative = "greater")
```

## Interpretation of the result



The **p-value** of the test is 0.01327, which is less than the significance level  $\alpha = 0.05$ . We can conclude that men's average weight is significantly different from women's average weight with a **p-value** = 0.01327.

## Access to the values returned by t.test() function

The result of **t.test()** function is a list containing the following components:

- **statistic**: the value of the **t test statistics**
- **parameter**: the **degrees of freedom** for the **t test statistics**
- **p.value**: the **p-value** for the test
- **conf.int**: a **confidence interval** for the mean appropriate to the specified **alternative hypothesis**.
- **estimate**: the means of the two groups being compared (in the case of **independent t test**) or difference in means (in the case of **paired t test**).

The format of the **R** code to use for getting these values is as follow:

```
# printing the p-value  
res$p.value
```



```
[1] 0.0132656
```

```
# printing the mean  
res$estimate
```


```
mean in group Man mean in group Woman  
68.98889          52.10000
```

```
# printing the confidence interval  
res$conf.int
```

```
[1] 4.029759 29.748019  
attr(,"conf.level")  
[1] 0.95
```

## Online unpaired two-samples t-test calculator

You can perform unpaired **two-samples t-test, online**, without any installation by clicking the following link:



[Online two-samples t-test calculator](#)

## See also

- Compare one-sample mean to a standard known mean
  - [One-Sample T-test \(parametric\)](#)
  - [One-Sample Wilcoxon Test \(non-parametric\)](#)
- Compare the means of two independent groups
  - [Unpaired Two-Samples Wilcoxon Test \(non-parametric\)](#)

## Infos



This analysis has been performed using **R software** (ver. 3.2.4).





Enjoyed this article? I'd be very grateful if you'd help it spread by emailing it to a friend, or sharing it on Twitter, Facebook or Linked In.

Show me some love with the like buttons below... Thank you and please don't forget to share and comment below!!

Share 11

Like 11

Tweet


Share

Save


Share

37


## Recommended for You!



Machine Learning Essentials:  
Practical Guide in R




Practical Guide to Cluster Analysis in  
R



Practical Guide to Principal  
Component Methods in R



R Graphics Essentials for Great Data Visualization



Network Analysis and Visualization in R



[More books on R and data science](#)

## Want to Learn More on R Programming and Data Science?

Follow us [by Email](#)

by [FeedBurner](#)

On Social Networks:  
on Social Networks



**Get involved :**



Click to **follow us** on [Facebook](#) and [Google+](#) :



**Comment this article** by clicking on "Discussion" button (top-right position of this page)

This page has been seen 188084 times

Newsletter



Boosted by [PHPBoost](#)

Recommended for you



Be Awesome in ggplot2:  
A Practical Guide to b...

[www.sthda.com](http://www.sthda.com)



Correlation matrix : An  
R function to do all yo...

[www.sthda.com](http://www.sthda.com)



GGPlot Cheat Sheet for  
Great Customization...

[www.sthda.com](http://www.sthda.com)



Be Awesome in ggplot2:  
A Practical Guide to b...

[www.sthda.com](http://www.sthda.com)

AddThis

