



[Lecture 16: Goodness of Fit Tests](#)

[Continued: Kolmogorov-Smirnov](#)

[test, Kolmogorov-Lilliefors test,](#)

3. The Empirical Cumulative

[Course](#) > [Unit 4 Hypothesis testing](#) > [Quantile-Quantile Plots](#)

> Distribution Functions

3. The Empirical Cumulative Distribution Functions

True or False on CDF

1/1 point (graded)

True or False: Let X be a random variable with cdf $F(x)$. Then

$$F(x) = P(X \leq x) = \mathbb{E}[\mathbf{1}(X \leq x)],$$

where $\mathbf{1}$ is the indicator function.

☒ True

☐ False



Solution:

By definition,

$$\begin{aligned} F(x) &= P(X \leq x) = \int_{-\infty}^x f(t) dt \\ &= \int_{-\infty}^{\infty} f(t) \mathbf{1}(X \leq x) dt \\ &= \mathbb{E}[\mathbf{1}(X \leq x)]. \end{aligned}$$

Submit

You have used 1 of 1 attempt

i Answers are displayed within the problem

Empirical Cumulative Distribution



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Empirical Cumulative Distribution Function

Let X be a random variable with distribution \mathbf{P} . Recall the cdf of \mathbf{P} is given by the function

$$F : \mathbb{R} \rightarrow [0, 1]$$

$$t \mapsto \mathbf{P}(X \leq t).$$

Let $X_1, \dots, X_n \stackrel{iid}{\sim} X$. The **empirical cumulative distribution function**, also called the **empirical cdf**, is the random function

$$F_n : \mathbb{R} \rightarrow [0, 1]$$

$$t \mapsto \frac{1}{n} \sum_{i=1}^n \mathbf{1}(X_i \leq t).$$

The empirical cdf depends on n and the observed data $X_i, i = 1, \dots, n$.

Example of Empirical CDF

5/5 points (graded)

Let X_1, \dots, X_5 be i.i.d. random variables. You obtain the sample $X_1 = 5, X_2 = 1.5, X_3 = -3, X_4 = 0.0, X_5 = 7$.

Let $F(t)$ be the empirical cdf of this sample. Find

Find $F(-4)$.

$F(-4) =$

✓ Answer: 0

Find $F(-3)$.

$F(-3) =$

✓ Answer: 1/5

Find $F(10)$.

$F(10) =$

✓ Answer: 1

Find the largest interval of t for which $F(t) = 3/5$. Answer by entering A and B in the equation below:

$F(t) = 3/5$ for $A \leq t < B$ where

$A =$ ✓ Answer: 1.5

$B =$ ✓ Answer: 5

Solution:

Given the sample $X_1 = 5, X_2 = 1.5, X_3 = -3, X_4 = 0.0, X_5 = 7$, the empirical cdf is

$$F(t) = \frac{1}{5} \sum_{i=1}^5 \mathbf{1}(X_i \leq t)$$
$$= \begin{cases} 0 & \text{if } t < -3 \\ 1/5 & \text{if } -3 \leq t < 0.0 \\ 2/5 & \text{if } 0 \leq t < 1.5 \\ 3/5 & \text{if } 1.5 \leq t < 5 \\ 4/5 & \text{if } 5 \leq t < 7 \\ 1 & \text{if } 7 \leq t \end{cases}.$$

Hence $F(-4) = 0, F(-3) = 1/5, F(10) = 1$, and $F(t) = 3/5$ for $1.5 \leq t < 5$.

Remark: The empirical cdf is right-continuous, just like the cdf.

Submit

You have used 1 of 3 attempts

❗ Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 4 Hypothesis testing:Lecture 16: Goodness of Fit Tests Continued: Kolmogorov-Smirnov test, Kolmogorov-Lilliefors test, Quantile-Quantile Plots / 3. The Empirical Cumulative Distribution Functions

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.



© All Rights Reserved