



[Course](#) > [Unit 1: Fourier Series](#) > [Period 2L](#) > 2. Recall Fourier series

Audit Access Expires Jun 24, 2020

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

2. Recall Fourier series

Recall that for any function $f(t)$ that is periodic and has period 2π , the Fourier series is given by

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nt + b_n \sin nt),$$

where the formulas for the coefficients a_n and b_n are given by

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \cos(nt) dt,$$


$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \sin(nt) dt$$



By virtue of the fact that we have formulas for the coefficients, a function has only one Fourier series. That is, if $f(t) = g(t)$, then the Fourier series for $f(t)$ is the same as the Fourier series for $g(t)$.

We will use this idea to come up with easier ways to compute Fourier series.

Here are the Fourier series of some 2π -periodic functions.

Name	Function	Fourier series	Graph
Square wave	$\text{Sq}(t) = \begin{cases} 1 & 0 < t < \pi \\ -1 & -\pi < t < 0 \end{cases}$	$\text{Sq}(t) = \frac{4}{\pi} \sum_{n \text{ odd}} \frac{\sin(nt)}{n}$	
Sawtooth wave	$W(t) = t, -\pi < t < \pi$	$W(t) = 2 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin(nt)$	
Triangle wave	$T(t) = t , -\pi < t < \pi$	$T(t) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n \text{ odd}} \frac{\cos(nt)}{n^2}$	

2. Recall Fourier series

Topic: Unit 1: Fourier Series / 2. Recall Fourier series

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.



[Learn About Verified Certificates](#)

© All Rights Reserved

