

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 2 Foundation of Inference](#) > [Lecture 5: Delta Method and Confidence Intervals](#)

7. Estimating the Parameter for an Exponential Model

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

7. Estimating the Parameter for an Exponential Model

Estimating the Parameter for an Exponential Model

guy?

Well, the variance of T_1 , which as you can tell from here,

is one over λ squared.

OK?


And that's two applications of integration by part.

Right?

You have to integrate T squared against e to the minus λT . OK?

And every time you're going to integrate,

you're going to have one λ that goes to the denominator.



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Consistency and Biasedness

4/4 points (graded)

Let $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \exp(\lambda)$. Let $\bar{X}_n := \frac{1}{n} \sum_{i=1}^n X_i$ denote the sample mean of the data set.

To which value does \bar{X}_n converge (both a.s. and in probability) as $n \rightarrow \infty$?
(Choose all that apply)

☒ $\mathbb{E}[X_i]$

☐ $\frac{1}{\mathbb{E}[X_i]}$

☐ $\mathbb{E}\left[\frac{1}{X_i}\right]$

☐ λ

☒ $\frac{1}{\lambda}$



To which value does $\frac{1}{\bar{X}_n}$ converge (both a.s. and in probability) as $n \rightarrow \infty$? (Choose all that apply)

☐ $\mathbb{E}[X_i]$

☒ $\frac{1}{\mathbb{E}[X_i]}$

☐ $\mathbb{E}\left[\frac{1}{X_i}\right]$

☒ λ

☐ $\frac{1}{\lambda}$



Which of the following is the bias of $\frac{1}{\bar{X}_n}$ as an estimator of λ ? (Choose all that apply.)

☒ $\mathbb{E} \left[\frac{1}{\bar{X}_n} \right] - \lambda$

☒ $\mathbb{E} \left[\frac{1}{\bar{X}_n} \right] - \frac{1}{\mathbb{E}[X_i]}$

☒ $\mathbb{E} \left[\frac{1}{\bar{X}_n} \right] - \frac{1}{\mathbb{E}[\bar{X}_n]}$

☐ $\frac{1}{\mathbb{E}[X_i]} - \lambda$

☐ $\frac{1}{\mathbb{E}[X_i]} - \frac{1}{\mathbb{E}[\bar{X}_n]}$



Which of the following are properties of $\frac{1}{\bar{X}_n}$ as an estimator of λ ? (Choose all that apply.)

☒ consistent

☐ unbiased

**Solution:**

- By the (strong/weak) law of large numbers

$$\bar{X}_n = \frac{\sum_{i=1}^n X_i}{n} \xrightarrow[n \rightarrow \infty]{a.s./\mathbf{P}} \mathbb{E}[X_i] = \frac{1}{\lambda}.$$

- On the other hand, by the continuous mapping theorem

$$\frac{1}{\bar{X}_n} \xrightarrow[n \rightarrow \infty]{a.s./\mathbf{P}} \frac{1}{\mathbb{E}[X_i]} = \lambda.$$

- Hence, we can answer the last part immediately: $\frac{1}{\bar{X}_n}$ is a consistent estimator of λ .

- However,

$$\mathbb{E}\left[\frac{1}{\bar{X}_n}\right] \neq \frac{1}{\mathbb{E}[\bar{X}_n]} = \lambda.$$

So the bias of $\frac{1}{\bar{X}_n}$ as an estimator of $\lambda = \frac{1}{\mathbb{E}[X_i]} = \frac{1}{\mathbb{E}[\bar{X}_n]}$ is

$$\text{Bias} = \mathbb{E}\left[\frac{1}{\bar{X}_n}\right] - \frac{1}{\mathbb{E}[\bar{X}_n]}.$$

Remark: Since the function $\frac{1}{x}$ is convex (by the shape of its graph or by $\left(\frac{1}{x}\right)'' = \frac{2}{x^3} > 0$), Jensen's inequality gives $\mathbb{E}\left[\frac{1}{\bar{X}_n}\right] > \frac{1}{\mathbb{E}[\bar{X}_n]}$

and hence the bias is greater than zero.

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Review: Central Limit Theorem

1/1 point (graded)

The **Central Limit Theorem** states that if X_1, \dots, X_n are i.i.d. and

$$\mathbb{E}[X_1] = \mu < \infty ; \quad \text{Var}(X_1) = \sigma^2 < \infty,$$

then

$$\sqrt{n} \left[\left(\frac{1}{n} \sum_{i=1}^n X_i \right) - \mu \right] \xrightarrow[n \rightarrow \infty]{(d)} Z \quad \text{where } Z \sim \mathcal{N}(0, ?).$$

What is $\text{Var}(Z)$? (Express your answer in terms of n , μ and σ).

$\text{Var}(Z) =$

sigma^2

✓ Answer: sigma^2

σ^2

STANDARD NOTATION

Solution:

For any n ,

$$\text{Var} \sqrt{n} (\bar{X}_n - \mu) = n \text{Var} (\bar{X}_n) = \text{Var} (X_i) = \sigma^2.$$

The central limit theorem states as $n \rightarrow \infty$, the distribution of $\sqrt{n} (\bar{X}_n - \mu)$ becomes Gaussian with the variance above (and mean 0); that is,

$$\sqrt{n} (\bar{X}_n - \mu) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, \sigma^2).$$

Note: The variance of Z is called the **asymptotic variance** of \bar{X}_n , even though it equals the variance of $\sqrt{n} \bar{X}_n$.

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 2 Foundation of Inference: Lecture 5: Delta Method and Confidence Intervals / 7. Estimating the Parameter for an Exponential Model

Add a Post

Show all posts ▼

by recent activity ▼

? Review: Central Limit Theorem

Could someone please elaborate on the **note** on **asymptotic variance** in the review problem's solution? I don't understand its significance.

2

💬 Poor Notation

Since Z is almost always used to represent a standard normal variable/distribution, I think another letter should be used here say W .

1

Learn About Verified Certificates

© All Rights Reserved