



[Course](#) > [Unit 1: Fourier Series](#) > [2. Properties of Fourier Series \(of Period 2L\)](#) > 6. Functions of arbitrary period


Audit Access Expires Jun 24, 2020

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

6. Functions of arbitrary period

Manipulating Fourier series



▶ 9:25 / 9:25

▶ 1.50x 🔊 🗑️ 📄 🗣️

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)


[Download Text \(.txt\) file](#)

Everything we did with periodic functions of period 2π can be generalized to periodic functions of other periods.

Problem 6.1 Define




$$f(t) := \begin{cases} 1 & \text{if } 0 < t < 1, \\ 0 & \text{if } -1 < t < 0. \end{cases}$$



and extend it to a periodic function of period 2. Express this new square wave $f(t)$ in terms of S_q . Then use the known Fourier series of S_q to find the Fourier series for $f(t)$.

Solution: This function is neither even nor odd. However, if we shift the function downwards by $1/2$, it is an odd function.




Write the shifted function


$$s(t) = f(t) - 1/2 = \begin{cases} 1/2 & 0 < t < 1 \\ -1/2 & -1 < t < 0 \end{cases}.$$

If we multiply this function by 2,


$$2s(t) = 2f(t) - 1 = \begin{cases} 1 & 0 < t < 1 \\ -1 & -1 < t < 0 \end{cases}.$$



what we get is very similar to the square wave of period 2π .



To avoid confusion, let's use u as the variable for Sq. Scaling the horizontal axis by factor of $1/\pi$ produces the graph of $2s(t)$.



In other words, if t and u are related by $u = \pi t$ (so that $u = \pi$ corresponds to $t = 1$), then $2s(t) = \text{Sq}(u)$. In other words,

$$2s(t) = \text{Sq}(\pi t).$$

Therefore,



$$f(t) = \frac{1}{2} + s(t) = \frac{1}{2} + \frac{1}{2} \text{Sq}(\pi t).$$

Since we know the Fourier series

$$\text{Sq}(u) = \frac{4}{\pi} \sum_{n \text{ odd}} \frac{\sin(nu)}{n},$$

the Fourier series for $f(t)$ is

$$f(t) = \frac{1}{2} + \frac{2}{\pi} \sum_{n \text{ odd}} \frac{\sin(n\pi t)}{n}.$$

Period $2L$ functions

Similarly we can scale the horizontal axis of any function of period 2π to get a function of different period. Let L be a positive real number. Start with "any" periodic function

$$g(u) = \frac{a_0}{2} + \sum_{n \geq 1} a_n \cos nu + \sum_{n \geq 1} b_n \sin nu,$$

of period 2π . Stretching horizontally by a factor L/π gives a periodic function $f(t)$ of period $2L$, and "every" f of period $2L$ arises this way. By the same calculation as above,

$$f(t) = g\left(\frac{\pi t}{L}\right)$$

$$= \frac{a_0}{2} + \sum_{n \geq 1} a_n \cos \frac{n\pi t}{L} + \sum_{n \geq 1} b_n \sin \frac{n\pi t}{L}$$



The substitution $u = \frac{\pi t}{L}$ (and $du = \frac{\pi}{L} dt$) also leads to Fourier coefficient formulas for period $2L$:

$$\begin{aligned}a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} g(u) \cos nu \, du \\&= \frac{1}{\pi} \int_{-L}^L g\left(\frac{\pi t}{L}\right) \cos\left(\frac{n\pi t}{L}\right) \frac{\pi}{L} dt \\&= \frac{1}{L} \int_{-L}^L f(t) \cos\left(\frac{n\pi t}{L}\right) dt.\end{aligned}$$

A similar formula gives b_n in terms of f .

6. Functions of arbitrary period

Topic: Unit 1: Fourier Series / 6. Functions of arbitrary period

[Hide Discussion](#)[Add a Post](#)

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

[Learn About Verified Certificates](#)

© All Rights Reserved

