



[Course](#) > [Unit 5 Bayesian statistics](#) > [Homework 9: Bayesian Statistics](#) > 4. Bayesian Estimation

4. Bayesian Estimation

Setup:

In this problem, we will explore the intersection of Bayesian and frequentist inference. Let $X_1, X_2, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} N(0, \theta)$, for some unknown positive number θ , which is our parameter of interest. Suppose that we are unable to come up with a prior distribution for θ .

(a)

1.0/1 point (graded)

Compute the maximum likelihood estimator of θ . You may use the variables n , $\sum_{i=1}^n X_i$, and $\sum_{i=1}^n X_i^2$.

(Enter **Sigma_i(X_i)** for $\sum_{i=1}^n X_i$ and **Sigma_i(X_i^2)** for $\sum_{i=1}^n X_i^2$. Do not worry if the parser does not render properly; the grader works independently. If you wish to have proper rendering, enclose **Sigma_i(X_i)** by brackets.)

$\hat{\theta}^{\text{MLE}} =$ ✓ Answer: Sigma_i(X_i^2)/n

[STANDARD NOTATION](#)

Solution:

Generating Speech Output

$$\hat{\theta} = S_n = \frac{1}{n} \sum_i (X_i)^2$$

Submit

You have used 2 of 3 attempts

i Answers are displayed within the problem

(b)

2/2 points (graded)

Is the MLE $\hat{\theta}^{\text{MLE}}$ asymptotically normal?

☒ It is asymptotically normal

☐ It is **not** asymptotically normal



If it is asymptotically normal, what is its asymptotic variance $V(\theta)$? If it is not asymptotically normal, type in 0.

$V(\theta) =$

2*theta^2

✓ Answer: 2*theta^2

2 · θ^2

STANDARD NOTATION

Solution:

We can construct L_n and l as following.

Generating Speech Output

$$L_n(x_1, \dots, x_n | \theta) = (2\pi\theta)^{-n/2} \exp\left\{-\sum_i \frac{1}{2\theta} x_i^2\right\}$$

$$l = \log L_1 = -\frac{1}{2} \log(2\pi\theta) - \frac{1}{2\theta} x_1^2$$

$$\frac{\partial l}{\partial \theta} = -\frac{1}{2\theta} + \frac{1}{2\theta^2} x_1^2$$

$$\frac{\partial^2 l}{\partial \theta^2} = \frac{1}{2\theta^2} - \frac{1}{\theta^3} x_1^2$$

$$I(\theta) = \frac{1}{2\theta^2}$$

Therefore, we can conclude that it is asymptotically normal, and its asymptotic variance is $2\theta^2$.

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, 2\theta^2)$$

Submit

You have used 3 of 3 attempts

i Answers are displayed within the problem

Generating Speech Output

(c)

2.0/2 points (graded)

Let's take a Bayesian approach here to arrive at an estimator.

Perform the following steps:

- Compute Jeffreys prior.
- Use Bayes formula to compute the posterior distribution.
- From the posterior distribution, compute the Bayesian estimator of θ . Recall that this is defined in lecture to be the mean of the distribution.

What is the Bayesian estimator $\hat{\theta}^{\text{Bayes}}$?

(Enter **Sigma_i(X_i)** for $\sum_{i=1}^n X_i$ and **Sigma_i(X_i^2)** for $\sum_{i=1}^n X_i^2$. Do not worry if the parser does not render properly; the grader works independently. If you wish to have proper rendering, enclose **Sigma_i(X_i)** and **Sigma_i(X_i^2)** by brackets.)

$\hat{\theta}^{\text{Bayes}} =$ ✓ Answer: Sigma_i(X_i^2)/(n-2)

In this Bayesian problem, which, if any, of the prior or the posterior, is proper?

☐ The prior only.

☒ The posterior only.

☐ Both the prior and the posterior.

☐ Neither the prior nor the posterior.



Solution:

1. Compute Jeffreys prior. Is it proper?

$$\pi_j(\theta) \propto \sqrt{\det(I(\theta))} = \frac{1}{\sqrt{2\theta}}$$

Since $\frac{1}{\sqrt{2\theta}}$ integrates to infinity, the prior is improper.

2. Use Bayes formula in order to compute the posterior distribution.

$$\pi(\theta|X_1, \dots, X_n) \propto \pi(\theta) L_n(x_1, \dots, x_n|\theta) \propto \frac{(2\pi\theta)^{-n/2}}{\sqrt{2\theta}} \exp\left\{-\sum_i \frac{x_i^2}{2\theta}\right\} \propto \theta^{-(n+2)/2} \exp\left\{-\sum_i \frac{x_i^2}{2\theta}\right\}$$

The posterior distribution is Inverse Gamma with parameters $\alpha = \frac{n}{2}, \beta = \frac{1}{2} \sum_i X_i^2$

3. Compute the Bayesian estimator of θ associated with Jeffreys prior.

$$\hat{\theta} = \int \theta \pi(\theta|X_1, \dots, X_n) d\theta = \frac{1/2 \sum X_i^2}{n/2 - 1} = \frac{\sum X_i^2}{n - 2}$$

Submit

You have used 2 of 3 attempts

i Answers are displayed within the problem

Generating Speech Output

0 points possible (ungraded)

Consider the set of statements given below. Decide whether each of the statements below is true. If a statement is true, decide whether this reflects a Bayesian or frequentist property of the Bayesian estimator.

Note: This problem is about the Bayesian estimator that you obtained in (c).

A **frequentist property** refers to all estimator properties that were considered before this lecture and are used in the context where there is a fixed, true, parameter value, and we want our estimator to approximate this value.

On the other hand, a **Bayesian property** refers to properties that indicate that we weight the likelihood somehow, not just using the raw values of the likelihood. (This could involve no additional judgement on the value of the parameter, for example if we use Jeffreys prior.)

Which statements are true and reflects a Bayesian property of the Bayesian estimator in (c)?

☐ The Bayesian estimator is consistent.

☐ The Bayesian estimator is unbiased.

☒ The Bayesian estimator gives in expectation a larger estimate if there are few observations, given a fixed θ , due to the nature of the prior used.

☐ The Bayesian estimator does not assume any particular prior distribution that is independent of the conditional likelihood.

☐ The Bayesian estimator is asymptotically normal.



Which statements are true and reflects a frequentist property of the Bayesian estimator in (c)?

☒ The Bayesian estimator is consistent. ✓

☐ The Bayesian estimator is unbiased.

Generating Speech Output

☐ The Bayesian estimator gives in expectation a larger estimate if there are few observations, given a fixed θ , due to the nature of the prior used.

☐ The Bayesian estimator does not assume any particular prior distribution that is independent of the conditional likelihood. ✓

☐ The Bayesian estimator is asymptotically normal. ✓

✗

Solution:

X_i is drawn from the distribution $N(0, \theta)$, so we compute

$$\mathbb{E}[X_i^2] = \mathbb{E}[X_i]^2 + \text{Var}(X_i) = 0^2 + \theta = \theta,$$

and

$$\mathbb{E}\left[\sum_{i=1}^n X_i^2\right] = n\theta.$$

- The Bayesian estimator is consistent because

$$\mathbb{E}[\hat{\theta}] = \frac{1}{n-2} \mathbb{E}\left[\sum_{i=1}^n X_i^2\right] = \frac{n}{n-2} \theta \rightarrow \theta$$

as $n \rightarrow \infty$, as

Generating Speech Output

$$\lim_{n \rightarrow \infty} \frac{n}{n-2} = 1.$$

Consistency is a frequentist property of an estimator as it is only desired when we assume a true value for the parameter.

- The Bayesian estimator is not unbiased, because as we calculated earlier,

$$\mathbb{E}[\hat{\theta}] = \frac{n}{n-2}\theta \neq \theta.$$

- The expected value of the Bayesian estimator is $\frac{n}{n-2}\theta$, which is decreasing in n , so it is true that this is larger when there are only a few observations. Indeed, this is due to the prior used, $\frac{1}{\sqrt{2\theta}}$, which gives larger weight to smaller values of θ . This is a Bayesian property because the notion of having our observations matter more strongly when we have more observations is strongly related to the Bayesian concept of starting from a prior distribution then updating through our observations.
- This Bayesian set-up uses the Jeffreys prior, which depends completely on the conditional likelihood and is thus a non-informative prior. Despite being a prior, this reflects a frequentist property of the procedure (and thus the estimator), because we do not assume any distribution beyond what's contained in the model and hence in some sense have no "prior". In fact, Jeffreys prior bridges the ideological gap between frequentist and Bayesian statistics.
- It is true that the Bayesian estimator is asymptotically normal because it converges in distribution to $\mathbb{E}[X_i^2]$ which is an expectation of an average, which is known to be asymptotically normal by the Central Limit Theorem. Asymptotic normality again is a property that's desired in the frequentist approach, because we want the parameter to have a predictable frequency distribution over the true parameter.

Submit

You have used 2 of 2 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Generating Speech Output

Bayesian statistics: Homework 9: Bayesian Statistics / 4. Bayesian Estimation

Show all posts ▼

by recent activity ▼

- ? [\[Staff\] Removal of Constant on Prior](#) 1
- 💬 [Solutions in Julia](#) 2
9.4 using SymPy @vars $\Sigma x^2 \theta n x$ # MLE for Normal(0, θ) $L_{akn} = (1/(\text{sympy.sqrt}(2*\text{sympy.pi} * \theta)^n)) * \exp(-\Sigma x^2 / (2*\theta))$ $lLn = \log(L_{akn})$ $dL1_ \theta = \text{simplify}(\text{diff}(lLn, \theta))$ $\text{solve}(dL1_ \theta...$
- ✓ (b): [Why the Fisher I matrix is equal to the expected hessian and NOT to its negative ?](#) 3
- ? [confused about the explanation of question \(d\)](#) 5
- 💬 [STAFF - Can you pls check my answer for part a](#) 3
[It seems like a straight forward formula based question that i got wrong. Can you pls check my answer and grader.](#)
- ? [STAFF - Wrong entry.](#) 3
[Hi, By error, I overwrite a good answer to question a \(I believed I was entering a value for C in fact\). Do you have a way to get my previous answer back ? BR, S.](#)
- 💬 [Hint: Read about the Inverse-Gamma Distribution](#) 16
[after hours of scratching, finally... hope this helps someone.](#)
- ? [Please help\(c\)](#) 3
[I have gotten Jeffrey's prior, multiplied that by the normal likelihood function and solved the gamma inverse parameters in order to get the mean of the gamma inverse for th...](#)
- 💬 [FYI: question \(d\) is ungraded. Staff, I recommend using more prominent text to communicate this](#) 3
[Staff, I recommend making this more clear. If one looks carefully one notices the 0/0 points - however, this text is not very prominent and I doubt most students read this for ...](#)
- ? [Part c\) MAP](#) 2
- ? [What's the intuition of having bias when using improper uniform prior in \(c\) instead?](#) 2 new_
[I'm not sure I understand the denominator being what it is \(n-4\). A lot of the time it seems like the prior \(for beta cases for example\) seems explainable in the posterior mean ...](#)
- 💬 [To those ones who are struggling with the last page exercises.](#) 2
[I'm adding this post because this particular topic was vaguely covered and I actually STRUGGLED A LOT until I found what to do. I INVITE EVERYONE TO ADD **NON-HOMEWO...](#)