

edX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#). ✕



[Lecture 3: Parametric Statistical](#)

[Course](#) > [Unit 2 Foundation of Inference](#) > [Models](#)

> 7. Examples of Parametric Models

**Audit Access Expires Dec 24, 2019**

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 7. Examples of Parametric Models

### Review: Sample Spaces of Distributions

4/4 points (graded)

Recall that a **sample space** of a random variable  $X$  is a set that contains all possible outcomes of  $X$ .

Note that the sample space of  $X$  is *not unique*. For example, if  $X \sim \text{Ber}(p)$ , then both  $\{0, 1\}$  and  $\mathbb{R}$  can serve as a sample space. However, in general, we associate a random variable with its smallest possible sample space (which would be  $\{0, 1\}$  if  $X \sim \text{Ber}(p)$ ).

Find the **smallest sample space** for each of the following random variables.

$X_1 \sim \text{Poiss}(\lambda)$ , a **Poisson** random variable with parameter  $\lambda$ :

☐  $\{0, 1\}$

☒  $\{x \in \mathbb{Z} : x \geq 0\}$  ✓

☐  $[0, \infty)$

☐  $(-\infty, \infty)$

$X_2 \sim \mathcal{N}(0, 1)$ , a **standard Gaussian (or normal)** random variable with mean 0 and variance 1:

☐  $\{0, 1\}$

☐  $\{x \in \mathbb{Z} : x \geq 0\}$

☐  $[0, \infty)$

☒  $(-\infty, \infty)$  ✓

$X_3 \sim \exp(\lambda)$ , an **exponential** random variable with parameter  $\lambda > 0$ :

☐  $\{0, 1\}$

☐  $\{x \in \mathbb{Z} : x \geq 0\}$

☒  $[0, \infty)$  ✓

☐  $(-\infty, \infty)$

$X_4 \sim \mathcal{I}(Y > 0)$  where  $Y$  is standard Gaussian and  $\mathcal{I}$  is the **indicator function**.

Recall the definition of the indicator function is:

$$\mathcal{I}(Y > 0) = \begin{cases} 1 & \text{if } Y > 0 \\ 0 & \text{if } Y \leq 0. \end{cases}$$

☒  $\{0, 1\}$  ✓

☐  $\{x \in \mathbb{Z} : x \geq 0\}$

☐  $[0, \infty)$ ☐  $(-\infty, \infty)$ **Solution:**

- A Poisson random variable is discrete and can take values on all non-negative integers.
- Gaussian random variables can take any real value.
- The Exponential distribution is continuous and is restricted to all non-negative real values.
- The final random variable is an indicator, so it must take values in  $\{0, 1\}$ . Note that  $X_4$  is in fact Bernoulli.

**Submit**

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Examples of parametric and nonparametric models

I'm sorry:

It has to be an integer.

It has to be an integer.

But for a Poisson, this is more precise.



It's always going to be the case.  
So just give-- always, thank you,  
always give the most possible precise  
answer.

I could actually write  $z$  here.

Don't do it.

I could write  $z$ .

All integers.

But it's certainly a superset of what I'm  
looking for.

But I should not, because I'm never going  
to get any negative numbers.

So if I know ahead of time I'm only  
going to get positive integers, let  
me just write a non-negative integers,  
natural integer.

And here I write the same thing.

So I'm going to have this thing



## Video

[Download video file](#)

## Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

## Statistical Model Definition Concept check

1/1 point (graded)

Which of the following is a statistical model?

- ☐  $\left(\{1\}, (\text{Ber}(p))_{p \in (0,1)}\right)$
- ☒  $\left(\{0, 1\}, (\text{Ber}(p))_{p \in (0.2, 0.4)}\right)$  ✓

☐ Both of the above

☐ None of the above

### Solution:

Solution in video below.

The set  $\{1\}$  is not the sample space of the distribution  $\text{Ber}(p)$ , so the first choice  $\left(\{1\}, (\text{Ber}(p))_{p \in (0,1)}\right)$  is not a statistical model. On the other hand,  $\left(\{0, 1\}, (\text{Ber}(p))_{p \in (0.2, 0.4)}\right)$  is a valid statistical model.

**Remark:** In the model  $\left(\{0, 1\}, (\text{Ber}(p))_{p \in (0.2, 0.4)}\right)$ , the parameter  $p$  is restricted to be in the interval  $(0.2, 0.4)$ . Such a restriction is perfectly valid, and can be useful for performing modeling tasks.

Submit

You have used 1 of 2 attempts

**i** Answers are displayed within the problem

## A Non-Example of a Statistical Model

0 points possible (ungraded)

(This problem is strictly pedagogical and is ungraded.)

Let  $\mathcal{U}([0, a])$  denote the uniform distribution on the interval  $[0, a]$ . Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{U}([0, a])$  for some unknown  $a > 0$ . Which one of the following is *not* a statistical model associated with this statistical experiment?

☒  $([0, a], (\mathcal{U}([0, a]))_{a>0})$  ✓

☐  $(\mathbb{R}_+, (\mathcal{U}([0, a]))_{a>0})$

☐ Neither choice above is a statistical model.

### Solution:

See video below.

The first choice  $([0, a], (\mathcal{U}([0, a]))_{a>0})$  is not a statistical model because the sample space, as written, depends on an unknown parameter  $a$ .

The second choice  $(\mathbb{R}_+, (\mathcal{U}([0, a]))_{a>0})$  is a statistical model because for any value of  $a$ , the random variables  $X_1, \dots, X_n$  will have sample space contained in the interval  $[0, \infty) = \mathbb{R}_+$ .

Submit

You have used 2 of 2 attempts

**i** Answers are displayed within the problem

## Worked example: Definition of Statistical model

[Start of transcript. Skip to the end.](#)

### Exercises

a) Which of the following is a statistical model?

1.  $(\{1\}, (\text{Ber}(p))_{p \in (0,1)})$
2.  $(\{0, 1\}, (\text{Ber}(p))_{p \in (0.2, 0.4)})$
3. Both 1 and 2
4. None of the above

b) Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{U}([0, a])$  for some unknown  $a > 0$ . Which one of the following is the associated statistical model?

1.  $([0, a], (\mathcal{U}([0, a]))_{a > 0})$
2.  $(\mathbb{R}_+, (\mathcal{U}([0, a]))_{a > 0})$
3.  $(\mathbb{R}, (\mathcal{U}([0, a]))_{a > 0})$
4. None of the above



10/61

OK, so which one is a statistical model?

OK, so either-- well there's clearly only one answer

that's valid here.

So which one do we have?

Who says one?

Actually, let's just go for it.

Is one a valid statistical model?

Who says yes?

Who says no?





## Video

[Download video file](#)

## Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

## Discussion





Hide Discussion

**Topic:** Unit 2 Foundation of Inference:Lecture 3: Parametric Statistical Models / 7. Examples of Parametric Models

Add a Post

Show all posts ▼

by recent activity ▼

-  [Need some more details on sample space](#) 2
- Hi all, in the second video prof. Philippe explains why a sample space cannot be defined in terms of an unknown paramater. But I cannot manage to...
-  [\[STAFF\] Grader is wrong on "A Non-Example of a Statistical Model"](#) 3
- Not a big deal, as the question is ungraded, but the grader accepts a wrong answer and rejects the correct one.<br>Note that the explanation in "Sh...
-  [\[Staff\] Typo in the Lecture Notes](#) 5
-  [Is 'sample space' the domain or subset of the range of a random variable?](#) 9

Learn About Verified Certificates

© All Rights Reserved