

<u>Help</u>

sandipan_dey 🗸

Next >

<u>Course</u> <u>Progress</u> <u>Dates</u> <u>Discussion</u> <u>Syllabus</u> <u>Outline</u> <u>laff routines</u> <u>Community</u>

★ Course / Week 11: Orthogonal Projection, Low Rank Approximation,... / 11.3 Orthonorm...

()



☐ Bookmark this page

Previous

⊞ Calculator

Week 11 due Dec 22, 2023 21:12 IST Completed

11.3.2 Orthonormal Vectors

At 2:11 in the first video, Robert writes square root of 2 as the length of the vector when it should be square root of 5. Please make a note of it!

Video 11.3.2 Part 1



▶ 2.0x

X

CC

66

Start of transcript. Skip to the end.

Dr. Robert van de Geijn: Let's first look at what orthonormal vectors are.

So here we're given k vectors.

Let's call them q0 through qk minus 1.

They're all vectors in Rm.

We're going to say that these vectors are mutually orthonormal.

If all i and j, the inner product of those two vectors is always 0.

Video

▲ Download video file

▶ 0:00 / 0:00

Transcripts

- ▲ Download SubRip (.srt) file
- ▲ Download Text (.txt) file

Reading Assignment

0 points possible (ungraded)
Read Unit 11.3.2 of the notes. [LINK]



Done



Submit

Discussion

Topic: Week 11 / 11.3.2

Hide Discussion

Add a Post

Show all posts

sts '

by recent activity \checkmark

There are no posts in this topic yet.

×

Homework 11.3.2.1

2/2 points (graded)

1. The vectors $\begin{pmatrix} -\sin{(\theta)} \\ \cos{(\theta)} \end{pmatrix}$, $\begin{pmatrix} \cos{(\theta)} \\ \sin{(\theta)} \end{pmatrix}$ are mutually orthonormal.

2. The vectors $\begin{pmatrix} \sin{(\theta)} \\ \cos{(\theta)} \end{pmatrix}$, $\begin{pmatrix} \cos{(\theta)} \\ -\sin{(\theta)} \end{pmatrix}$ are mutually orthonormal. True/False

Submit

Video 11.3.2 Part 2

TRUE



Start of transcript. Skip to the end.

Dr. Robert van de Geijn: So the answer turns out to be yes.

First of all, if you take the dot product of these two vectors,

what do you get?

You get minus the sine of theta, times the cosine of theta,

plus the cosine of theta, times the sine of theta.

Video

▲ Download video file

Transcripts

- <u>♣ Download SubRip (.srt) file</u>
- ▲ Download Text (.txt) file

Homework 11.3.2.2

1/1 point (graded)

Let $q_0,q_1,\ldots,q_{k-1}\in\mathbb{R}^m$ be a set of orthonormal vectors and let $Q=\left(egin{array}{c|c}q_0&q_1&\cdots&q_{k-1}\end{array}
ight)$.

Then $oldsymbol{Q}^Toldsymbol{Q}=oldsymbol{I}.$

TRUE ✓ Answer: TRUE

$$egin{array}{ll} egin{array}{ll} Q^T \ Q \end{array} &= \left(egin{array}{c|c} q_0 & q_1 & \cdots & q_{k-1} \end{array}
ight)^T \left(egin{array}{c|c} q_0 & q_1 & \cdots & q_{k-1} \end{array}
ight) = \ \left(egin{array}{c|c} q_0^T \end{array}
ight) \end{array}$$

$$egin{aligned} &=& \left[egin{array}{c|c|c|c} q_1^T \ \hline draw{d}_1^T \ \hline draw{d}_{k-1}^T \end{array}
ight] \left(egin{array}{c|c|c|c} q_0 & q_1^T q_1 & \cdots & q_0^T q_{k-1} \ \hline q_1^T q_0 & q_1^T q_1 & \cdots & q_1^T q_{k-1} \ \hline draw{d}_1^T q_0 & q_1^T q_1 & \cdots & q_1^T q_{k-1} \ \hline draw{d}_{k-1}^T q_0 & q_{k-1}^T q_1 & \cdots & q_{k-1}^T q_{k-1} \ \hline \end{array}
ight) = \left(egin{array}{c|c|c} 1 & 0 & \cdots & 0 \ \hline 0 & 1 & \cdots & 0 \ \hline draw{d} & draw{d} & draw{d} & draw{d} \ \hline draw{d} & draw{d} & draw{d} & draw{d} & draw{d} \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 0 \ \hline 0 & 1 & \cdots & 0 \ \hline draw{d} & draw{d} & draw{d} \ \hline \end{matrix}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline 0 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 0 \ \hline 0 & 1 & \cdots & 0 \ \hline draw{d} & draw{d} & draw{d} \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 1 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 0 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 0 \ \hline \end{array}
ight) \ &= \left(egin{array}{c|c|c} 1 & 0 & \cdots & 0 \ \hline \end{array}
ight) \ &= \left(eg$$

Submit

Answers are displayed within the problem

Video 11.3.2 Part 3



Start of transcript. Skip to the end.

Robert van de Geijn: And the answer is that this is true.


How do we show that?

Just go through it.

If you multiply Q transpose times Q,

the same as exposing the columns of Q in each of these matrices.

If you then transpose the first one you





Video

Download video file

Transcripts

Homework 11.3.2.3

1/1 point (graded)

Let $Q \in \mathbb{R}^{m imes k}$ (with $k \leq m$) and $Q^T Q = I$. Partition

$$Q = \left(egin{array}{c|c} q_0 & q_1 & \cdots & q_{k-1} \end{array}
ight).$$

Then $q_0, q_1, \ldots, q_{k-1}$ are mutually orthonormal vectors.

TRUE

✓ Answer: TRUE

$$egin{aligned} egin{aligned} egi$$

Hence
$$q_i^Tq_j=egin{cases} 1 & ext{if } i=j \ 0 & ext{otherwise}. \end{cases}$$

Submit

Answers are displayed within the problem

Video 11.3.2 Part 4



Start of transcript. Skip to the end.

Dr. Robert van de Geijn: And the answer is that this is true.

How do we know that?

Well, if you look at Q transpose Q, you

that, which is then equal to that, which is then

equal to that, which is then equal to

Previous

Next >

© All Rights Reserved



edX

About

Affiliates

edX for Business

Open edX

Careers

<u>News</u>

Legal

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

<u>Trademark Policy</u>

<u>Sitemap</u>

Cookie Policy

Your Privacy Choices

Connect

<u>Idea Hub</u>

Contact Us

Help Center

<u>Security</u>

Media Kit















© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>