

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 3 Methods of Estimation](#) > [Lecture 8: Distance measures between distributions](#)

15. Likelihood of a Poisson Statistical
> Model

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

15. Likelihood of a Poisson Statistical Model

Review: Statistical Model for a Poisson Distribution

2/2 points (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Poiss}(\lambda^*)$ for some unknown $\lambda^* \in (0, \infty)$. Let $(E, \{\text{Poiss}(\lambda)\}_{\lambda \in \Theta})$ denote the corresponding statistical model. What is the smallest possible set that could be E ?

☐ $\mathbb{N} = \{1, 2, 3, \dots\}$

☒ $\mathbb{N} \cup \{0\}$

☐ \mathbb{Z}

☐ \mathbb{R}



The parameter space Θ can be written as an interval (a, ∞) . What is the smallest value of a so that $\{\text{Poiss}(\lambda)\}_{\lambda \in (a, \infty)}$ represents all possible Poisson distributions?

$a =$

✓ Answer: 0.0

Solution:


A Poisson random variable takes values on all non-negative integers $\{0, 1, 2, \dots\}$. Hence, the smallest possible sample space is $\mathbb{N} \cup \{0\}$.

A Poisson random variable is specified by its mean λ , which is allowed to be any positive real number. Hence, $a = 0$ is the correct choice.

You have used 1 of 2 attempts

i Answers are displayed within the problem

Likelihood of a Poisson Statistical Model



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Practice: Compute Likelihood of a Poisson Statistical Model

3/3 points (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Pois}(\lambda^*)$ for some unknown $\lambda^* \in (0, \infty)$. You construct the associated statistical model $(E, \{\text{Pois}(\lambda)\}_{\lambda \in \Theta})$ where E and Θ are defined as in the answers to the previous question.

Suppose you observe two samples $X_1 = 1, X_2 = 2$. What is $L_2(1, 2, \lambda)$? Express your answer in terms of λ .

$$L_2(1, 2, \lambda) = e^{-(2\lambda)} \lambda^{3/2}$$

✓ Answer: $e^{-(2\lambda)} \lambda^{3/2}$

$$\frac{e^{-2\lambda} \lambda^3}{2}$$

Next, you observe a third sample $X_3 = 3$ that follows $X_1 = 1$ and $X_2 = 2$. What is $L_3(1, 2, 3, \lambda)$?

$$L_3(1, 2, 3, \lambda) = e^{-(3\lambda)} \lambda^{6/12}$$

✓ Answer: $e^{-(3\lambda)} \lambda^{6/12}$

$$\frac{e^{-3\lambda} \lambda^6}{12}$$

Suppose your data arrives in a different order: $X_1 = 2, X_2 = 3, X_3 = 1$. What is $L_3(2, 3, 1, \lambda)$?

$$L_3(2, 3, 1, \lambda) = e^{-(3\lambda)} \lambda^{6/12}$$

✓ Answer: $e^{-(3\lambda)} \lambda^{6/12}$

$$\frac{e^{-3\lambda} \lambda^6}{12}$$

STANDARD NOTATION

Solution:

The probability mass function of $\text{Pois}(\lambda)$ is $x \mapsto e^{-\lambda} \frac{\lambda^x}{x!}$ where $x \in \mathbb{N} \cup \{0\}$. Hence by definition

$$L_n(x_1, \dots, x_n, \lambda) = \prod_{i=1}^n e^{-\lambda} \frac{\lambda^{x_i}}{x_i!} = e^{-n\lambda} \frac{\lambda^{\sum_{i=1}^n x_i}}{x_1! \cdots x_n!}.$$

Hence, first we plug in $n = 2$, $x_1 = 1$, and $x_2 = 2$:

$$L_2(1, 2, \lambda) = e^{-2\lambda} \frac{\lambda^{1+2}}{2!1!} = e^{-2\lambda} \frac{\lambda^3}{2}.$$

When the next sample arrives, we can simply evaluate the density of a Poisson at the observation:

$$P(X_3 = 3) = e^{-\lambda} \frac{\lambda^3}{3!}, \quad X \sim \text{Pois}(\lambda)$$

and multiply this by the previous response:

$$L_3(1, 2, 3, \lambda) = e^{-\lambda} \frac{\lambda^3}{3!} L_2(1, 2, \lambda) = e^{-3\lambda} \frac{\lambda^6}{12}.$$

Remark 1: Observe that we can compute the likelihood sequentially as the data arrives, updating it in the previous fashion after each new observation.

Similarly, we see that

$$L_3(2, 3, 1, \lambda) = e^{-3\lambda} \frac{\lambda^6}{12}.$$

Remark 2: Observe that the likelihood of observations $X_1 = x_1, \dots, X_n = x_n$ is independent of the *order* in which these observations arrive.

You have used 1 of 3 attempts

Submit

 Answers are displayed within the problem

Properties of the Likelihood

1/1 point (graded)

Let $(E, \{P_\theta\}_{\theta \in \Theta})$ denote a discrete statistical model. Let p_θ denote the pmf of P_θ . Let $X_1, \dots, X_n \stackrel{iid}{\sim} P_{\theta^*}$ where the parameter θ^* is unknown. Then the **likelihood** is the function

$$L_n : E^n \times \Theta \rightarrow \mathbb{R}$$

$$(x_1, \dots, x_n, \theta) \mapsto \prod_{i=1}^n p_\theta(x_i).$$

For our purposes, we think of x_1, \dots, x_n as observations of the random variables X_1, \dots, X_n .

Which of the following are properties of the likelihood L_n ? (Choose all that apply.)

Hint: It may be useful to consider your responses from the previous question.

☐ The likelihood does not change with the parameter θ .

☒ The likelihood can be updated sequentially as new samples are observed. For example, $L_3(x_1, x_2, x_3, \theta) = L_1(x_3, \theta) L_2(x_1, x_2, \theta)$

☒ The likelihood is symmetric: it doesn't matter the order in which we plug in the observations. For example, $L_4(x_1, x_2, x_3, x_4, \theta) = L_4(x_2, x_3, x_1, x_4, \theta)$ and this is true for any rearrangement of x_1, x_2, x_3, x_4 .

☐ If we eliminate a single observation, then the likelihood remains unchanged. For example, $L_3(x_1, x_2, x_3, \theta) = L_2(x_1, x_2, \theta)$



Solution:

We examine the choices in order.

- "The likelihood does not change with the parameter θ ." is incorrect. Rather, it is crucial that we interpret the likelihood L_n as a function of θ . That is, L_n varies as θ ranges over the parameter space Θ . This is evident in the likelihoods for the Bernoulli and Poisson models in the previous problems.
- "The likelihood can be updated sequentially as new samples are observed. For example, $L_3(x_1, x_2, x_3, \theta) = L_1(x_3, \theta) L_2(x_1, x_2, \theta)$." is also correct. In the previous problem, we saw that to compute the likelihood after observing $X_3 = 3$, we simply took the old likelihood $L_2(1, 2, \lambda)$ and multiplied it by $L_1(3, \lambda)$. Note that $L_1(x_3, \theta) = p_\theta(x_3)$, the density of P_θ evaluated at the new observation. Inspection of the defining formula

$$L_n(x_1, \dots, x_n, \theta) = \prod_{i=1}^n p_\theta(x_i)$$

implies that the likelihood can be updated sequentially in this fashion.

- "The likelihood is symmetric..." is correct. We observed in the previous problem that observing the samples in a different order does not affect the likelihood. This is also evident from the definition of the likelihood: we can take the product


$$\prod_{i=1}^n p_\theta(x_i)$$

in any order, and the result will still be the same.

- "If we eliminate a single observation, then the likelihood remains unchanged..." is incorrect. In the previous question, we saw that for a Poisson statistical model, $L_2(1, 2, \lambda)$ and $L_3(1, 2, 3, \lambda)$ do not have the same formula. Hence, deleting an observation from the sample will change the likelihood.

Submit

You have used 1 of 2 attempts

 Answers are displayed within the problem

Discussion

[Hide Discussion](#)

Topic: Unit 3 Methods of Estimation:Lecture 8: Distance measures between distributions / 15. Likelihood of a Poisson Statistical Model

[Add a Post](#)

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

[Learn About Verified Certificates](#)

© All Rights Reserved