

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#). ✕



[Course](#) > [Unit 2 Foundation of Inference](#) > [Lecture 4: Parametric Estimation and Confidence Intervals](#)

2. Statistics, Estimators, Consistency,  
> and Asymptotic Normality

### Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)


## 2. Statistics, Estimators, Consistency, and Asymptotic Normality

### Statistics, Estimators, Consistency, and Asymptotic Normality

which is 0 almost all the time.

So I decide to use the word "asymptotic variance"

to denote the variance of  $\hat{\theta}_n$ ,  
but once rescaled by the square root of  $n$ ,



so that I have something that's meaningful

and not always equal to 0.

**Is that clear?**

[End of transcript. Skip to the start.](#)



## Video

[Download video file](#)

## Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

# Which Statistics are Estimators?

1/1 point (graded)

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} P_\theta$  where the distribution  $P_\theta$  depends on an unknown parameter  $\theta \in \mathbb{R}$ . Which of the following statistics are considered **estimators**?

(Choose all that apply.)

☐  $\theta$

☒ 4.2

☒  $\sum_{i=1}^n i^2 X_i^i$

☒  $\frac{1}{n} \sum_{i=1}^n X_i$

☐  $\frac{1}{n} \sum_{i=1}^n X_i - \theta$



**Solution:**

Recall that heuristically, a statistic is a function of the data that can be easily computable, and an estimator is a statistic whose expression does not depend on the unknown parameter  $\theta$ .

Note that the first and last choices,  $\theta$  and  $\frac{1}{n} \sum_{i=1}^n X_i - \theta$  both have some explicit dependence of  $\theta$ , so they cannot be estimators.

On the other hand, the remaining expressions 4.2,  $\sum_{i=1}^n i^2 X_i^i$ , and  $\frac{1}{n} \sum_{i=1}^n X_i$  only depend on  $X_1, \dots, X_n$  (and not  $\theta$ ), so they are indeed estimators.

**Remark:** The second estimator, 4.2, is potentially a very poor choice as it does not depend on the data set. But according to the definition, it is still considered an estimator.

Submit

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Consistency of an Estimator

1/1 point (graded)

An estimator  $\hat{\theta}_n$  is **weakly consistent** if  $\lim_{n \rightarrow \infty} \hat{\theta}_n = \theta$ , where the convergence is in probability.

Suppose that in the previous problem the unknown parameter  $\theta$  is the common mean of  $X_1, \dots, X_n$ . Assume that  $\theta \neq 4.2$ . Which of the following is a weakly consistent estimator for  $\theta$ ? (Choose all that apply.)

☐  $\theta$

☐ 4.2☐  $\sum_{i=1}^n i^2 X_i$ ☒  $\frac{1}{n} \sum_{i=1}^n X_i$ ☐  $\frac{1}{n} \sum_{i=1}^n X_i - \theta$ **Solution:**

By the weak law of large numbers,  $\frac{1}{n} \sum_{i=1}^n X_i \rightarrow \mathbb{E}[X_1] = \theta$  in probability, so this is the correct choice.

From the previous question, the first and last choice,  $\theta$  and  $\frac{1}{n} \sum_{i=1}^n X_i - \theta$ , are not even estimators, so these options are incorrect.

Since  $\theta \neq 4.2$ , this estimator cannot be consistent. Finally, there is no guarantee that  $\sum_{i=1}^n i^2 X_i$  converges to  $\theta$ . In fact, for many choices of distribution, this statistic will diverge to  $\infty$ .

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Quantifying Consistency

1/1 point (graded)

**Note:** The problem statement has been changed to asking about convergence in probability instead of almost surely. Attempts will be reset.

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Ber}(p)$ . Let  $\bar{X}_n$  be the estimator given by  $\frac{1}{n} \sum_{i=1}^n X_i$ .

What is the smallest constant  $c$  such that

$$n^c \left( \bar{X}_n - p \right) = n^c \left( \frac{1}{n} \sum_{i=1}^n X_i - p \right)$$

does **not** converge to 0 in probability as  $n \rightarrow \infty$ ?

✓ Answer: .5

**Solution:**

Let  $\sigma = \sqrt{p(1-p)}$  denote the common standard deviation of  $X_1, \dots, X_n$ . By the central limit theorem,

$$\frac{\sqrt{n}}{\sigma}(\bar{X}_n - p) = \frac{\sqrt{n}}{\sigma} \left( \frac{1}{n} \sum_{i=1}^n X_i - p \right) \rightarrow N(0, 1)$$

where the convergence is in distribution. As a result, we see that for  $c < 1/2$ ,

$$n^c (\bar{X}_n - p) = \frac{\sigma}{n^{1/2-c}} \frac{\sqrt{n}}{\sigma} (\bar{X}_n - p) \approx \frac{\sigma}{n^{1/2-c}} N(0, 1) \rightarrow 0$$

in probability as  $n \rightarrow \infty$ . Hence,  $c = 1/2$  is the smallest possible value of  $c$  such that

$$n^c (\bar{X}_n - p) = n^c \left( \frac{1}{n} \sum_{i=1}^n X_i - p \right)$$

does **not** converge to 0 in probability as  $n \rightarrow \infty$ .

**Remark:** As defined in the third video in this section, this implies that the estimator  $\bar{X}_n$  is  $\sqrt{n}$ -consistent. This means that the estimator  $\bar{X}_n$  converges to the true parameter at a relatively fast rate, so this gives us something stronger than just consistency.

Submit

You have used 2 of 3 attempts

 Answers are displayed within the problem

## Discussion

[Hide Discussion](#)

**Topic:** Unit 2 Foundation of Inference:Lecture 4: Parametric Estimation and Confidence Intervals  
/ 2. Statistics, Estimators, Consistency, and Asymptotic Normality

[Add a Post](#)

### ◀ All Posts

## Problem#3: Quantifying Consistency

question posted 5 days ago by [rickytyagi](#)

I'm not following the logic in the solution, someone kindly clarify.

This post is visible to everyone.



### **Jeevesh1**

3 days ago - marked as answer 3 days ago by [rickytyagi](#)



Let  $X$  denote the expression in question without the  $n^c$  outside.

1.) Whenever the sample mean converges a.s. to  $p$ ,  $X$  will converge a.s. to 0 .

2.) Sample mean converges to  $p$  a.s (By strong Law of Large Numbers)

3.)If you multiply the expression  $X$  by  $\sqrt{n}$  , you will get a Standard Normal Gaussian.( $Z$ ). Now we see what would have happened , if we would have used a different power of  $n$  ??



4.) If you now divide  $Z$  by  $n^c$ ,  $0 < c < 0.5$  your gaussian will go to 0 at every point as  $n \rightarrow \infty$ . This is equivalent to multiplying  $X$  by some power of  $n$  less than 0.5 .

5.) If you would use some power of  $n$ , higher than 0.5, it'd be equivalent to multiplying the Gaussian  $Z$  , by some positive power of  $n$ . Which would cause the Gaussian to explode at all points as  $n \rightarrow \infty$  .

Show Comments (8) ▼

Preview

Submit

Learn About Verified Certificates

© All Rights Reserved