




Course > Section... > 1.3 A M... > 1.3.2 Q...

1.3.2 Quiz: Making Sense of the Lambert-Beer Model

🔖 Bookmark this page

Here's the Lambert-Beer model Margo presented:



[View Larger Image](#)

[Image Description](#)

μ uniform attenuation coefficient (units cm^{-1})

I_0 input x-ray intensity (units kVp)

I output x-ray intensity (units kVp)

Δx length of object (units cm)

$$I = I_0 e^{-\mu \Delta x}$$

Units of the Lambert Beer Model

- The units of intensity are **peak kilovoltage**, kVp, which is the maximum voltage applied across an X-ray tube.

- The units of Δx , the thickness, are centimeters, or cm.
- What are the units of μ , the attenuation coefficient? They are determined by the other units. Because $I = I_0 e^{-\mu \Delta x}$ must have units kVp, and I_0 has units kVp as well, this means we want this means we want $e^{-\mu \Delta x}$ to be unitless. So the units of μ must cancel those of Δx , and hence the units of μ are 1/cm or cm^{-1} .

Question 1

1/1 point (graded)

From the Lambert-Beer model, we know that the intensity of an x-ray of initial intensity I_0 passing through an object with attenuation coefficient μ and length Δx is given by

$$I = I_0 e^{-\mu \Delta x}$$

Compare the resulting intensities of light when a x-ray with initial intensity I_0 passes through each of the following objects.

Choose all that are correct.

Attenuation and Thickness Combinations

A: $\mu = 0.5, \Delta x = 1$

B: $\mu = 0.5, \Delta x = 2$

C: $\mu = 1, \Delta x = 2$

D: $\mu = 2, \Delta x = 0.5$

E: $\mu = 3, \Delta x = 3$

☐ E results in the largest output intensity

☒ A results in the largest output intensity ✓

☐ Output intensity for C > Output intensity for B

☒ Output intensity for B is the same as the Output intensity for D. ✓

☐ None of the above.



Explanation

The intensity is determined by the product of the thickness and attenuation coefficient. Because $f(t) = e^{-t}$ is always decreasing, smaller values of $t = \mu \cdot \Delta x$ correspond to larger values of I . The value of $\mu \cdot \Delta x$ is smallest in A and equal for B and C.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Question 2

1/1 point (graded)

Margo explained that attenuation depends on both the material (measured by the attenuation coefficient μ) and the thickness of the material. According to the Lambert-Beer model, do objects that attenuate more light (because of material or thickness or both) result in lesser output intensity? (Why or why not?)

☒ Yes, always. ✓

☐ Sometimes, but not always.

☐ Never.

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)



English ▼

© 2012–2018 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open edX logos are registered trademarks or trademarks of edX Inc. | 粤ICP备17044299号-2

POWERED BY
OPENedX®

